AMERICAN JOURNAL OF PHARMACY

AND THE SCIENCES SUPPORTING PUBLIC HEALTH

Since 1825

COMMITTEE ON PUBLICATION

† Charles H. LaWall, Ph. M., Sc. D. Mitchell Bernstein, M. D. J. W. Sturmer, D. Sc. John K. Thum, Ph. M. § Arno Viehoever, Ph. D. E. Fullerton Cook, Ph. M.

IVOR GRIFFITH, Ph. M., Sc. D., Editor Linwood F. Tice, M. Sc., Assistant to the Editor John E. Kramer, B. Sc., Business Manager

Vol. 109.

DECEMBER, 1937

No. 12

CONTENTS

Editorial:	
Words, Words	586
Original Articles:	
Field Tests for Marihuana (Cannabis). By Arno Viehoever, Phila- delphia, Pa.	589
Some Little Bug Is Going to Get You! By T. Swann Harding, Washington, D. C.	594
Abstracts from, and Reviews of, the Literature of the Sciences Supporting Public Health.	601
Solid Extracts.	615
Index to Volume 109.	619

† Deceased

Annual Subscription, \$3.00 Foreign Postage, 25 Cents Extra Single Numbers, 30 Cents. Back Numbers, 50 Cents

> Eniered as Second-Class Matter at the Post Office at Philadelphia, Pa., Under the Act of March 3, 1879

Acceptance for Mailing at Special Rate of Postage Provided for in Section 1103, Act of October 3, 1917. Authorized February 15, 1920

EDITORIAL

On these pages the editor offers his opinions, unshackled by advertising patrons and unrestrained by anything save a sense of the decent and the truthful—the editor, alone, is responsible for their type, their tone and their tenor.

WORDS, WORDS, WORDS

M OST of the world's misunderstandings come because of an ignorance of the value of words or because of a wilful or witless misconception of their intent and arrangement. There are words that weep and words that laugh, words of learned length, winged words and burning words, airy words and heavy words, words, true

and false, brave and cringing, sweet and sour-words, words, words.

The male American of average intelligence is said to have at his disposal a collection of some seven thousand words,—more than there is in the Old Testament,—but strangely enough the female of the indigenous species has but five thousand at her command.

But what a command!

And what a rapid turnover of stock!

The origin of words within a spoken language mirrors better than most men know, the course of history and the curve and verve of evolution.

Whence came our simplest words?

Whence, indeed, came first, the word word?

Consider the way in which English words, of which there are now, according to Max Müller, the celebrated philologist, some 100,000, have changed since Chaucer's day, and through various causes, chiefly because of invasion or invention. In Britain, since the dim Druidic days, layers after layers of words cover the primitive island tongue, as certainly as the seventeen soils of Ur hide the Abrahamic substrate. In the living English tongue of today the initiated may find some British residues. But much easier is it to find the words that came with Caesar's legions and the Danes and the Anglo-Saxons, those flaxen-haired fighters from north of the North Sea. Easier still to find are the verbal increments of William the Conqueror, whose Norman hosts refurnished the isle with romantic actions as well as Romantic words.

Yes, indeed, the English dictionary, especially a good one, is rich in the miniatures of history for those who search aright, and also rich in sheer interest. Consider, for instance, the devious way in which we came to possess in an vocabulary the pretty word, *sincere*.

When Italian vase makers wished to give semblance of perfection to a cracked or blemished vase, the fault was filled with a wax and covered with paint.

Sine cera (without wax) was the term accordingly applied to a perfect vase, a genuine article. Hence and thence, our word sincere.

Abundance, in a wave (from unda, wave).

Alarm, which now suggests clockwise, "out of bed in a hurry," was once a "call to arms" to meet impending danger.

Alphabet, from the Greek a and b, alpha, beta.

Anecdote, from the Greek anekdotos, which means "not to be published," in which is a gentle hint to those who run their anecdotes to print.

Bank, from an origin which places the Banks of Loch Lomond in the same category with the Federal Reserve Banks, both incidentally being inhabited chiefly by Scots, and the word, as does "bench," coming from a Latin origin meaning a table, or elevation.

Bless and Blood have an odd association, in that blessing once meant "to consecrate by blood."

Bombastic, applied to speech, actually means stuffed with cotton (bombax).

Calculate, from calx or calculus, a stone, comes to us from the days when the Romans did their counting with stones or beads on a string.

Curfew literally means cover the fire and go to bed (couvre-feu, Fr.).

Symposium, to drink together (potio, to drink), now, of course, suggesting to think together.

And so might we continue, in merry analytic mood, except for the pedagogic fact that there is a better gain in doing this alone.

Let the reader then do his research in words, in a self-conducted fashion. Such research registers more fully.

And let this be the undersigned's contribution to the search, beyond just the mere suggestion. Here, in addition to the Webster, which still orates and perorates for those who listen, is a list of books where the study of word origins may be profitably conducted.

Anderson-New Study of English Words.

Greenough and Kittredge-The Words and Ways of English Speech.

Marsh-Origin and History of the English Language.

Skeat-English and Iceland.

Trench-Study of Words.

White-Words and Their Uses.

Whitney-Linguistic Science.

Picturesque Word Argus-The Merriam Company.

IVOR GRIFFITH.

Antiseptics and Wound Healing. W. W. Saeger, E. B. Vedder and C. Rosenberg, Am. J. Surg. 38, 348 (1937). Artificial wounds made in healthy dogs were infected with staphylococcus aureus. One wound in each dog was treated with a different antiseptic, the other was not treated and was used as a control. Tincture of metaphen and merthiolate, azochloramide and a 2 per cent. aqueous gentian violet solution were used on several dogs. The infected wounds were not improved by any of these antiseptics, due to injury of the cells caused by them.

L. G.

Copper Sulphate Treatment of Trichophytosis. J. B. Moloney, U. S. Naval Med. Bull. 35, 440 (1937): Fifty cases of trichophytosis (athlete's foot) gave complete cures by daily application of a 20 per cent. solution of dehydrated copper sulfate in glycerin (c. p.). The copper salt was dehydrated by heating over a Bunsen burner to a constant weight and care was taken to preserve the finished mixture from moisture. This was rubbed thoroughly into the affected area by inunction for fifteen minutes every day. The excess was removed and a dry sterile dressing was applied. This preparation is recommended as an alternative treatment in refractory cases.

L. G.

ORIGINAL ARTICLES

FIELD TESTS FOR MARIHUANA (Cannabis)*†

By Arno Viehoever

Marihuana, the dangerous habit-forming drug has latterly spread its tentacles, and accelerated its pace, in growth and harm, in our country. Here is a test to determine, quickly, its presence in cigarettes.

THE rapid spread in the illegal sale and use of the Mexican weed "Marihuana" in this country as a dope for chewing, or the smoking of cigarettes makes the need for a quick, simple, positive test, permitting its detection in waste lots, in the field or street an obvious fact.

As there are apparently no combined chemical and biological field tests now available to the narcotic enforcement officials or the analysts called upon to identify away from their laboratory with fair reliability the presence of marihuana in a cigarette or of the weed wherever grown or found, we have tried to devise such tests.

From a literature survey of the physical, chemical and biological data of cannabis and its physiologically active resin cannabinol, and an experimental check-up, the following field tests have been devised.

Chemical Field Test

A portion, one-third or more of a suspected cigarette (without the wrapper), or as little as one-tenth of a gram of the suspected weed, dried and somewhat comminuted, is placed in small, wide-mouthed containers. To it is added, followed by immediate shaking for five seconds, a small but excessive amount in volume of a test solution, consisting of benzene (nine parts) and sodium hydroxide (2 per cent.) dissolved in ethyl alcohol (one part). The liquid is then immediately decanted into an evaporating cup. Very satisfactory results have been obtained when cannabis was extracted with benzene for fifteen to ninety seconds, the slightly yellowish benzene solution then immediately decanted and mixed with equal amounts of alcoholic sodium hydroxide (2 per cent.).

Rapid removal of the solvent prevents the excessive extraction of green and yellow plant pigments (chlorophyll, etc.) which, more

^{*}A preliminary note from the Gross Laboratory for Biological and Biochemical Research, Philadelphia College of Pharmacy and Science.

[†] Presented before the A. A. A. S., Indianapolis, Ind., Dec. 27, 1937.

TABLE I
CHEMICAL FIELD TEST FOR CANNABIS

	Color Change	20 sec.	25 " Reagent was added di-	28 " rectly to material.	***	30 "	55 "	ı min.	86 sec.	85 " Benzene was added to	2 min. material and then de-	2 " Canted into INACII.	* 8	3 %	3½ min.	
Time	Decanta-	\$ sec.	* 20	5 "	2 20	2 20	3 20	" OI	3	, 6	3 %	ະ	3	3 6	,, 4	
	Extraction	5 sec.	3 10	,, 9	2 %	, 06	50 "	,, 09	20 "	* **	25 "	30 "	40 "	" OI	, OI	
	2% Alc. NaOH.					2 cc.		3 3	2 2		3		22 22	27 27	, , ,	
Amount	Benzene					2 cc.	2 2		" "	" "		29 29	3 3	. 29 29	20 20	
Ame	Reagent	3 cc.*	3 cc.*	3 cc.*	3 cc.**											
	Drug	0.1 gm.	3	27	22 23	3 3	n, n	33 33	" "	27 . 27	, ,					
	-1	-	a	3	4	מו	9	7	00	6	IO	11	122	13	14	

*1.5 cc. of a solution containing 1 cc. of 2% alc. NaOH + 4 cc. benzene, and 1.5 cc. benzene. ***1.5 cc. of a solution containing 1 cc. of 4% alc. NaOH + 4 cc. benzene, and 1.5 cc. benzene. ***Delayed color change, but observed within 24 hrs. due to high concentration of alc. NaOH.

or less, interfere with the speedy and unmistakable color development of the test.

A positive test, recognized by the appearance of a color change from slightly yellowish to pinkish within one to two minutes, becoming deeper red after standing a short time, indicated the presence of cannabinol. For its further characterization, spontaneous evaporation of the reddish liquid leaves a dry, partially pinkish to violet, rather persistent, residue. This dissolves to an orange-red solution in strong ammonia and with a violet to almost bluish-violet color in acetone.

Thus we believe to have worked out a field test, which is (1) simple, (2) quick to carry out, (3) reliable (when carried out as indicated above—with access of air and in the absence of an excess of water and carbonates, (4) conspicuous—visible by day and by lamplight, (5) delicate and (6) fairly permanent in liquid, the permanency increasing in the dry residue.

Biological Field Test

For a quick biological field test the residue from the benzene extraction, using 0.1 grant or more of the drug, is freed from benzene by the expediency of blowing on the liquid—to speed its complete evaporation—from a shallow glass cup.

The residue, by trituration is suspended in 2 cc. culture water, or more in case more drug is used. This culture water is poured on daphnia in V-tubes and the narcotic effects noted, which in the above concentration become strikingly noticeable within one hour. There is not only a marked slowing down of the locomotion of the animal but an obvious lowered activity of other essential body organs—a progressive toxic effect leading to death.

The narcotic effect may be depicted by "locomographs," obtained by direct tracing, per fifteen seconds time interval, of both the travel line of the normal daphnia and that of a narcotised animal. These locomographs, described more fully in a later contribution clearly show a progressive reduction in speed and distance of travel, then a lowered level of swimming and, finally, stationary locomotion in the advanced stages of narcosis.

Delayed cold, citalife, but boses see

The work, inclusive of the further study of the mechanism of action of cannabis and cannabinol by means of daphnia and other animals, is being continued with the assistance of Mr. Whiteley, working with a grant from the Alumni Association of the Philadelphia College of Pharmacy and Science.

SOME LITTLE BUG IS GOING TO GET YOU!

By T. Swann Harding, Washington, D. C.

A recent article, by the same writer, on the role of chemicals in disease, found so wide a range of appreciative readers, that the editor is pleased to follow it with this presentation in kindred vein.

YEARS ago they sang a song that ran "Some little bug is going to get you—someday!" After that the understanding was you passed on to your reward or punishment. Just about that time the present writer's mother was trundling him around among a lot of children who had mumps.

The idea was that he should go ahead and "catch" the mumps and be done with it. But he just wouldn't take the mumps. Other kids might do as they pleased but no mumps for him. So he passed them up. Then suddenly, spectacularly, and entirely unaccountably he came down with mumps when he was thirty-five, a disaster of which he refuses to speak further.

But why wouldn't he catch the mumps at nine? Where did he get them at thirty-five? Why did he have them then? One can look very wise and say that at nine he had a "natural immunity" to mumps and at thirty-five he did not. It often makes us feel very learned if we can translate our ignorance into polysyllables.

What is natural immunity? Science is not sure. Why will an individual frequently put up a powerful fight and conquer a virulent infection and then suddenly up and die from a minor one during

S

is

g

st

th

ge

tra

convalescence? Science says that is indeed baffling.

Of course there is such a thing as natural immunity. We just do not yet know what it is. Certain Algerian sheep are naturally immune to anthrax. Certain Zebu cattle are immune to dread footand-mouth disease which would wipe out the cattle industry in this country but for the alert United States Bureau of Animal Industry. Yet foot-and-mouth disease is mild, not deadly, in many other lands.

The Dutch elm disease, the chestnut blight, and the corn borer all raise hob in the United States, yet they do little or no serious damage in other regions of the world. Have the elms and chestnuts of Europe, for instance, developed a heritable form of immunity to these diseases which bid fair to wipe out our trees of these varieties?

Some animals seem to have germ-killing blood, or at least their blood carries in it antibodies or antitoxins which can counteract the chemical poisons produced by disease germs. Thus some horses as well as some children have naturally in their blood the antitoxin that combats diphtheria. Therefore they do not contract the disease.

If medicine could just discover the mechanism of this natural immunity and how it is inherited many diseases might be conquered. Advertising has, of course, made us infection conscious. Many are quite neurotic on the subject and take all sorts of absurd precautions to prevent infection.

Yet a wide variety of antiseptics are used before operations by different surgeons in different hospitals all over the land, or by different doctors on wounds. But the mortality is little affected by the nature of the antiseptic used. So much depends upon the individual patient's natural resistance to the particular germ and upon the virulence of the particular strain of germs, and their number, that it makes relatively little difference what antiseptic procedure is used. In some cases no antiseptic at all is almost surely best.

Again, we often read or hear about the importance of keeping the fingers out of the mouth and of cleansing the oral cavity. But there is no effective mouth and throat antiseptic. Moreover the upper part of the digestive tract contains a natural disinfecting agent and the stomach secretes strong acid. Germs that are swallowed are rapidly and quietly killed.

The lower part of the tract is populated by a wide variety of all sorts of germs. Down there things are alkaline instead of acid as is the case up above. The acid line largely determines how high the germs may rise in the tract. If fatigue or nervousness decreases the stomach acid, germs may begin to live in the stomach itself.

If rabbits swallow cholera germs while they are in normal health they do not contract cholera. But if the acid contents of their stomachs and upper intestines are first rendered alkaline, then the germs swallowed will kill them.

Since the stomach contents tend to be less acid in hot than in cool weather typhoid and dysentery are especially prevalent in summer. A diet which will maintain the stomach and upper intestinal tract in a state of normal acidity acts to ward off germ diseases.

Our clean skin is also a good germicide. It is a better germ killer than most antiseptics. The secretions of our mucous membranes, as well as the saliva, and gastric and intestinal juices all kill germs. We also have within us blood cells called leucocytes that eat germs alive.

The disease immunity of an individual is not the same at all times. Certain factors affect it. There is a wide range of individuasusceptibility to any germ disease. Then consider the health of the germs themselves. Some strains of germs seem to be weak, ill, or

lazy. They are not virulent any more.

An individual may, it appears, get used to certain germs of his own and carry them around constantly. They may look just like very dangerous germs, but they are not dangerous to him. But let him become infected with similar germs of a different strain. Then the resistance he has built up may break down and he may even lose his life.

Temperature also is a factor with which to reckon. A relatively low temperature will protect a lizard from the plague which he contracts readily when kept a little warmer. Birds are immune to the tuberculosis of cows, distinct from the bird tuberculosis from which they suffer, because their body temperature differs so much from that of a cow.

C

p it

S

S d

b

ir

th

It has long been held that fatigued and undernourished individuals easily contracted infections. But when experimental animals were compelled to exercise until fatigued and were then exposed to infection they did not prove more susceptible than unexercised controls. Studies of human beings working hard so as to induce muscular fatigue did not produce evidence of lowered resistance to infections.

Of course fatigue may have a different effect upon well-trained than upon untrained individuals. The effect of fatigue as a single factor seems hard to isolate. Certainly fatigue does not decrease the antibody content of the blood which determines resistance. But exhausting work immediately following an infection seems to produce drastic results.

Muscular fatigue has not been proved, however, to affect the body defense mechanism against infection one way or the other. Its influence appears to be minor and is difficult to reveal. The manner of infection, individual differences in susceptibility, and the varied virulence of the germs involved again assume more importance.

For a time we heard much about the anti-infective action of vitamin A and occasionally of other vitamins. The effect of vitamin B on immunity remains a mystery. Lack of vitamin C will lower a guinea pig's resistance to certain infections. There is little or no evidence to show that lack of vitamin D predisposes an animal to infections.

A deficiency of vitamin A does produce some loss of resistance in frection in rats, mice and guinea pigs. It may reduce resistance in early life in human beings. But it is not justifiable to call vitamin A "the anti-infective vitamin." Too many other factors may be involved. The normal antibody content of the blood serum and the power of the serum to produce these antibodies seem unaffected by dietary deficiencies.

Scurvy does lower a guinea pig's resistance to infection, hence the administration of vitamin C would be doubly helpful. Birds on a diet deficient in vitamin B are susceptible to anthrax in a way that normal birds are not. But respiratory infections appear to be just about as frequent among well-fed as among deficiently fed infants.

The difference between a liability to become infected and a capacity successfully to fight off an infection after it is contracted must always be borne in mind. Pronounced deficiencies in nutrition may lower resistance to infections, but few studies have been made of partial dietary deficiency—the kind most likely to occur in man—and its effects.

There is no evidence to show that increased feedings with any specific food element—protein, vitamin, mineral or what not—will increase the resistance of normal animals. Results obtained with one species of animals do not apply to other species either. Multiple dietary deficiencies of different kinds are common in man and may be important, but little attention had been paid in animal experiments to the kind of chronic lowered resistance that clinicians often find in children.

There is some evidence that infants are protected from certain infections by diets high in caloric content, vitamins and minerals, but even sex differences in resistance to infection, which are marked, often render such reports meaningless. There is some evidence also that many of the tubercular long lived on deficient diets.

Pigeons when suffering from a deficiency of vitamin B lose their natural resistance to anthrax. Dogs also contract certain infections when low in this vitamin which causes beriberi and pellagra in human beings. Yet sufferers from these diseases show no increased liability to infection.

An adequate diet before the onset of infection is far more important than the administration of vitamin A after infection sets in. On the other hand it has been shown that well-nourished rats are much more easily infected with foot-and-mouth disease than are ill-fed rats, while there are certain kinds of mouse tumors that will grow only on well-nourished and never on poorly nourished mice.

Cattle also seem to be more susceptible to foot-and-mouth disease when in top nutritional condition than when undernourished. Yet the acquired immunity to mouse typhoid is seriously impaired by starvation. Even if there is food in the stomach of a mouse an hour or so before it is infected, that does not help if it has previously been starved.

Underfed pigs also are more open to infestation with parasites than are the well fed. But this seems to be because the poorly fed animals are more likely to eat incautiously and get hold of food containing various worms and worm eggs than are the well-fed ones.

The germ itself also must be considered. It has to eat and live. It has a nutrition problem too. Germs require specific foods. The reason that human beings only are affected with gonorrhea is that the gonococci require a certain protein in their food that is found only in human beings.

Again meat-eating animals are seldom troubled with certain protozoa that infest the intestines of vegetarians. Animal proteins prevent the infestation. Carbohydrates intensify it. Casein will clear the protozoa out. Human beings can be cleared of these protozoa and cured of distressing symptoms after a week on a meat diet.

More and more evidence accumulates to show the chemical nature of disease. Germs injure us mostly because they produce chemicals that poison us. The viruses, which cause a long list of plant, animal and human diseases actually appear to be non-living chemical substances.

ir

SI

di

di

Why not attack diseases by chemical means then? This is what Ehrlich tried to do and the recent success of prontosil against certain streptococcic infections represents a further advance of his idea. Though germs are composed of protoplasm quite as human beings are, Ehrlich hoped to develop various chemicals, usually dyes, that would kills germs and not injure human tissues.

This was all based on the theory that diseases were caused chemically and that proteins had what he called "side-chains." The curative chemical must be able to hitch on to these side-chains in the body and put the deadly protein out of business as a disease cause. This theory worked in the case of syphilis and has other possibilities.

Another approach is by means of vaccines and antitoxins. These usually produce a temporary immunity. But some hold that resistance to smallpox has been built up through the years by all but universal vaccination. Today smallpox is milder even among the unvaccinated than it was formerly. Was a resistance built up and then inherited?

Sheep born of parents that were immunized against anthrax are said to have greater resistance to this disease than other sheep. A distinct resistance to certain serious diseases has also been built up in mice and in guinea pigs.

That brings up an important point. There is a good deal of evidence to indicate that resistance to certain diseases is hereditary. Agriculturalists have been breeding disease-resistant plants for many years. But such resistance is always specific for a certain disease and a certain type of organism.

A sugarcane bred resistant to one disease may prove susceptible to another and be wiped out. Or, again, if one strain of the organism causing the disease is baffled by that sugarcane it may hybridize, and a new strain may appear that will attack it.

Disease resistance has also been bred in animals, but it is specific, not generalized. Lines of mice resistant or susceptible to mouse typhoid have been bred. But not all laboratory mice originally show the same susceptibility to typhoid.

The germs are given them and a line of mice bred from the survivors shows surprising resistance to the disease. Whereas thirty-seven out of a hundred of the original mice died, only fifteen of the hundred will die in this resistant strain under the same conditions of infection.

Another line may be bred from the offspring of the mice which succumbed to the typhoid infection. Very soon a line will appear in which eighty-five to as high as ninety-seven out of every hundred will die when given mouse typhoid. But this resistance is specific. It is not associated with general bodily vigor. Nor will breeding mice of exceptional general vigor improve their resistance to a specific disease.

Nor is it ever possible to breed a wholly susceptible or a wholly resistant strain no matter how long this goes on. Like the writer and his mumps, some mice in the most susceptible line will refuse to take typhoid. So also, no matter how bad the plagues of the past have been, some always resisted, others survived the infection though attacked.

There does seem to be such a thing as species immunity. Hog cholera seems limited to hogs. Man is alone susceptible to a long list of diseases. Under usual conditions it is impossible to infect warm-blooded animals with diseases of the cold-blooded, and vice versa.

The same holds for insects. The larva of the rhinoceros beetle is very susceptible to cholera but is immune to anthrax and diphtheria. Crickets readily contract anthrax but never take bird tuberculosis. The anopheles mosquito, well-known carrier of malaria, is very susceptible to that disease while closely related types of mosquito are immune.

Does this apply also to human beings? When syphilis first appeared in Europe it was a raging and deadly plague. Today it is a comparatively mild disease among the so-called white race which seems relatively immune. It is a pest to other races.

Of course there are scientifically no races of men. There are only sub-species. When we say that the negro is more susceptible to tuberculosis than the white man we are also on dubious ground. In general the former have living conditions far inferior to the latter and that may account for the apparent difference in susceptibility.

B

d

g

bi

di

tu

pr

BI

be wl

COI

Even then negroes do appear to be relatively immune to diseases of the skin, to scarlet fever, diphtheria, measles, erysipelas, chickenpox, and to infections of the nasal and sinus regions. But as if to compensate that are markedly prone to such infections of the respiratory tract as pneumonia, bronchitis, and tuberculosis. Some attribute these differences to differences in tissue structure.

The Japanese are said to be immune to scarlet fever. Smallpox is a relatively mild disease in Mexico, though it was a dread scourge among the North American Indians. Negroes are believed immune to yellow fever. The Eskimo is immune to many diseases in his native habitat which, however, he readily contracts when he visits warmer climates.

Arabians are said to be very susceptible to tuberculosis when they live in cities. New York Russians, Poles and Jews appear to resist this disease better than do the Irish, though no race is concerned here, strictly speaking. Measles is mild among Caucasians but a severe plague among American Indians and Melanesians. The Chinese are relatively immune to lockjaw. The Malaysians are said to be more susceptible to beriberi than any other people.

But beriberi is a nutritional or a deficiency disease, so that proves only that the Malaysians can stand less denial of vitamin B than some other people, if it proves anything. Really there is not a great deal to prove. On the other hand it is much more important for us to know the limitations of our knowledge than to be certain about so many things that are not so at all. Then, at least, we know in which direction we should progress.

Reverting now to the original question of why the little mumps bug refused to nip the writer when he was nine but flattened him out when he was thirty-five—medical science does not yet know the answer. We have seen herein that only a partial answer is possible. Maybe the bugs were not very virulent one time. Possibly the individual was immune to the specific bug type at the time. Perhaps the bugs were more numerous on one occasion than upon the other.

The whole secret is probably locked in the chemical storeroom. Both germs and tissues are largely composed of proteins. Certain disease-producing agents that we formerly thought alive, but which are now regarded as non-living—the viruses—can, if they get upon the right tissue, assume some of the functions of life. They can grow, reproduce and mutate. Yet they are not actually living things.

Proteins are made up of twenty or more simpler chemicals called amino acids. These form chains. Specific amino acids can be combined in a wide variety of ways; they can take their places at many different points in the complex chains that make up proteins. The possible number of proteins is therefore almost infinite. The figure would at least be astronomical.

In other words it is far from impossible that the protein structure of each individual differs subtly and ever so slightly from the protein structure of any other individual. Bloods are known to type. Blood donors must have blood which types thus and so before it can be transfused into a patient. The proteins of different varieties of wheat have already been shown to differ slightly in their amino acid composition.

May not disease resistance be bound up in that complex protein molecule, a molecule so heavy and complex that it may at times have an atomic weight of half a million or even more? Is it not possible that an infective virus can hitch on to the side chain of only certain types of proteins and that the proteins of individuals differ? Is it not just as possible that the proteins of the same individual differ at different times in life?

Germs are composed in great part of the basic substance composing all living matter, the complex called protoplasm. This, in turn, largely consists of protein. Germs also have to eat, as we have seen. They are particular about their food also, as we have seen. Man has certain diseases because his proteins are the only kind that certain germs really relish.

Then what may we conclude? Natural immunity and disease infection are explicable chemically. The secret is locked within the living cell. It is the secret of protein constitution, a secret that can be learned only when we have much more refined methods than we have today.

The whole science of medicine is tending towards chemistry and away from absolute dependence on bacteriology. Recent work with the minute filtrable viruses causing smallpox, foot-and-mouth disease, infantile paralysis, possibly influenza, and many other ills, has shown that these ultramicroscopic disease-producing agents are proteins. But they become quasi-living on certain tissues.

Plants, animals and human beings whose proteins are of a certain composition resist these viruses; those with proteins of another general type are susceptible. Do we begin to see light here? It seems that we do but science warrants no more positive statement as yet.

ABSTRACTS FROM AND REVIEWS OF THE LITERATURE OF THE SCIENCES SUPPORTING PUBLIC HEALTH

Bacteriology						I	oui	is	Ger	shenfeld, B. Sc., Ph M.
Biochemistry	7,	Nut	riti	on,	ef	c.				Arno Viehoever, Ph. D.
Biology .										Marin S. Dunn, Ph. D.
										. Arthur Osol, Ph. D.
										Fullerton Cook, Ph. M. and their assistants

Treatment of Psoriasis With Massive Doses of Crystalline Vitamin D and Irradiated Ergosterol. E. T. Ceder and L. Zon, Public Health Reports 52, 1580 (1937). Psoriasis is recognized as one of the most frequent disorders of the skin. The numerous concepts regarding its nature have led to the use of a variety of therapeutic agents only a few of which have been found to afford any benefit to the majority of cases.

In the past decade viosterol in small doses has been employed in the treatment of psoriasis without signal success although Krafka (J. Lab. and Clin. Med. 21, 1147 (1936)) reported benefit following larger doses of viosterol in three cases. Only in recent years have the more potent irradiated ergosterol products been available and been found to be apparently safe. The unfortunate results in the past are said by Bills to have been due to the presence of toxisterol which is now absent in commercial products. Furthermore, the reports in the literature show that the deleterious effect of massive doses of irradiated ergosterol have occurred in children and in rats which have since been shown to be much more sensitive than the adults of either species.

The authors while employing large doses of vitamin D averaging 300,000 units daily in the treatment of chronic arthritis observed a complete involution of a coexisting psoriasis. This led to the use of this substance in the treatment of fifteen cases of chronic widespread psoriasis. Of the total number eleven obtained a complete involution in a maximum of twelve weeks, while two obtained only a partial benefit and two showed no benefit. The results indicated furthermore that such massive doses appear to be relatively safe, inasmuch as in the few instances of hypervitaminosis produced, no extreme symptoms developed. The study surely indicates great promise in the treatment of psoriasis by this method.

L. F. T.

Scoops as a Source of Contamination of Ice Cream in Retail Stores. A. J. Krog and D. S. Dougherty, Amer. J. Pub. Health 27, 1007 (1937). A comparative study of the results of routine ice cream analyses conducted at Plainfield, N. J., over several years showed that the bacterial counts of loose ice cream were in most instances considerably higher than those of the same product manufactured and packaged at the same plant and dispensed by the same dealer.

A comparison study of the bacterial count of loose ice cream collected with the vendor's scoop and with a sterile spoon was made with the following results:

Samples taken with vendor's scoop	Samples taken with sterile spoon
640,000	8,000
320,000	23,000
512,000	3,000
158,000	28,000
121,000	3,000
113,000	10,000
19,000	2,000
175,000	11,000

Further examinations led to the following interesting conclusions:

1. The amount of ice cream in the can has a direct effect on the count since the freshly opened can would not have been subjected to the high degree of scoop contamination as a nearly empty can.

2. The time of the day samples were taken influenced the count due to the fact that dippers and containers were usually washed in the morning after the establishment was opened.

3. The physical condition of the scoop was important, those mended with plugs of cloth and those never properly washed giving higher contamination. The usual arrangement of keeping scoops in a container of water is objected to as it provides in the course of use an excellent medium for the rapid multiplication of bacteria. A new method of keeping scoops by placing them in a dry rack protected from flies, dust, etc., and washing them in running water both before and after use was found to greatly lower the bacterial content of ice cream dipped in this manner. Ice cream retailers should be interested in this worthwhile recommendation.

L. F. T.

The Isolation of Crystalline Vitamin A. H. N. Holmes and R. E. Corbett, J. A. C. S. 59, 2042 (1937). The authors report the isolation of crystals of vitamin A from three different samples of fish liver oil. The crystals when crystallized from methanol at very low temperature appear in beautiful rosettes or radiating clusters. They are optically inactive and isotropic, melting point 7.5-8.0 degrees. The extinction coefficient ($E_{1_{\text{em}}}^{16}$) was found by Darby of Columbia to be equal to 2100 and by biological assay Dutcher and Guerrant of Pennsylvania State College found it to contain considerably above 2,265,000 and somewhat below 3,400,000 international units per gram. The molecular weight, determined by the freezing point lowering of cyclohexane, was found to be about 294. This is less than 3 per cent. above the weight corresponding to the formula $C_{20}H_{30}O$. The difference is explained as probably due to the rapid change in the structure of the vitamin in solution through oxidation.

L. F. T.

Safeguards Proposed to Govern Distribution of Dangerous Drugs. In a report submitted to Congress by the Secretary of Agriculture as a result of the recent deaths resulting from the use of elixir of sulfanilimide, four recommendations are made as follows:

I. License control of new drugs to insure that they will not be generally distributed until experimental and clinical tests have shown them to be safe for use. The definition of what constitutes a new drug should include (a) substances which have not been used sufficiently as drugs to become generally recognized as safe, (b) combinations of well-known drug substances where such combinations

have not become generally recognized as safe, and (c) well-known drug substances and drug combinations bearing label directions for higher dosage or more frequent dosage or for longer duration of use than has become generally recognized as safe. Exemption should be made for new drugs distributed to competent investigators for experimental work. A board of experts should be provided who will advise the Secretary of Agriculture on the safety of new drugs.

2. Prohibition of drugs which are dangerous to health when administered in accordance with the manufacturer's directions for

3. Requirement that drug labels bear appropriate directions for use and warnings against probable misuse.

4. Prohibition of secret remedies by requiring that labels disclose fully the composition of drugs. Many foreign countries now impose this requirement. Many drugs manufactured in the United States are exported to such countries under labels bearing such disclosure. The same drugs are sold to our citizens under labels that give no hint of their composition.

L. F. T.

The Stabilization of Adrenalin Solutions. B. Sjögren and H. Larsson, Farmacevtisk Revy 309 (1936), through Pharmas. Zentrh. 78, 695 (1937). The question of the stabilization of adrenalin solution was studied and sodium metabisulfite demonstrated the best stabilizing effect of all the inorganic reducing agents. An adrenalin solution when properly prepared with this substance is stable almost indefinitely. The sodium metabisulfite is, even in the concentration of 0.1 per cent., quite efficient and it is likewise of value since it is only very slightly toxic, the adrenalin is in no way chemically altered, nor is its pharmacological or physiological action.

L. F. T.

Signs of Vitamin A Deficiency in the Eye Correlated With Urinary Lithiasis. W. J. Ezickson and J. B. Feldman, J. A. M. A. 109, 1706 (1937). The authors investigated any possible relationship between vitamin A deficiency and upper urinary lithiasis in human beings. A group of individuals who had or did have renal or

te

sl

te

th

uretal calculi were tested for vitamin A deficiency by the dark adaption or light sensitivity test. Of twenty-five patients with urolithiasis, twenty-four were found to have pathologic dark adaption varying from mild to severe and investigations disclosed dietary deficiency in vitamin A in many instances. The patients were treated with doses of vitamin A ranging from 13,000 to 52,000 units daily and, in addition, they were placed on acid ash or alkaline ash diets depending on the urinary pH. The results were in accord with the clinical studies reported by the Council on Pharmacy and Chemistry of the American Medical Association, namely, that vitamin A fails to correct dark adaption where it exists together with urolithiasis. By this study it seems evident that the vitamin A deficiency responsible for lack of dark adaption occurs in association with renal lithiasis. but such deficiency is dependent on lack of assimilation or utilization of this substance rather than on dietary deficiency. The possibility that lack of vitamin A assimilation or utilization and urinary lithiasis may have a common metabolic basis is suggested.

Vitamin A Content of Cod Liver Oil. A Comparison of Spectrophotometric and Chemical Methods. A. D. Holmes, F. Tripp and G. H. Satterfield, J. Ind. and Eng. Chem., Analytical Edition 9, 456-57 (1937). Thirty-two samples of cod liver oil have been examined for their vitamin A potency, using the Hilger vitameter E value and the antimony trichloride blue value methods. A comparison of the results obtained by the two methods indicated that in general they were of the same order. The oils that gave high E values also gave high blue values, while those with the lowest E values also gave the lowest blue values. The ratio blue value to E value for most samples was between nine to one and eleven to one.

The results obtained for free fatty acid and unsaponifiable material showed that the amount of these materials present in an oil is not correlated with its E value or blue value; no consistent relationship has been found between vitamin A potency of an oil and its content of free fatty acid or unsaponifiable material.

L. A. R.

Nicotine Thiocyanate—A Contact Insecticide. J. S. McHargue and R. K. Calfee, J. Ind. Eng. Chem. 29, 1232 (1937). Nicotine thiocyanate was prepared by decomposition of ammonium thio-

cyanate. The compound is found to be very effective as an aphidicide and, in sufficient concentration and with a suitable spreader, controls red spider. It is decomposed by the more positive ions in solution, losing much of its effectiveness, and is then capable of serious foliage injury. Practical spreaders for use with this insecticide are discussed.

L. J. K.

Synthetic Aliphatic Penetrants. B. G. Wilkes and J. N. Wickert, J. Ind. and Eng. Chem. 29, 1234 (1937). During the past few years an increasing interest in the use of compounds possessing surface activity has taken place. Among such compounds the sodium secondary alcohol sulfates have recently become available to industry in quantity. The chemical structure of these and a number of other compounds possessing surface activity are described. A comparison is made of their relative wetting action, surface tension and lime soap dispersibility under neutral, acid and alkaline conditions. In addition the authors describe some commercial applications in which these sodium secondary alcohol sulfates are or can be used to advantage.

L. J. K.

Sterilization of Aqueous Solutions of Sodium Phenobarbitone. H. W. Tomski and L. J. Waller, Pharm. J. 139, 3859 (1937). The instability of aqueous solutions of sodium phenobarbitone having been confirmed by Nielsen (1933), the authors carried out an investigation to determine the loss of the drug when different methods of sterilization were applied to a solution. The loss was estimated by the volumetric method of Hegland (1935) in which sodium carbonate solution is added to the barbiturate solution, the resulting mixture being titrated with N/10 AgNO3 until the faint opalescence produced does not disappear on shaking. Experiments were also made to determine the loss of sodium phenobarbitone when N/10 sodium carbonate is added and the resulting solution subsequently sterilized. In the sterilizing of a 20 per cent. solution of sodium phenobarbitone, using the various methods of sterilization, the following results were obtained: By autoclaving, 115 degrees C. for thirty minutes, loss of 19 to 20.5 per cent.; heating at 80 degrees C., under the process of tyndallization, loss of 6 to 7 per cent.; boiling for fifteen minutes, loss of I per cent.; heating at 100 degrees C. for one hour, loss of 10 per cent. When subjected to any of the above methods of sterilization the 20 per cent, solutions formed a white

precipitate of phenylethylacetylurea on cooling.

From the experiments the authors recommend, solutions of sodium phenobarbitone for injection should be prepared from a sample yielding a solution of a pH value as low as possible and should be sterilized by filtration only. The solutions should be stored only for a short period and in a cool place. H. P. F.

Bacterial Filtration. H. Berry, Pharm. J. 139, 3853, 3854 (1937). The sterilization of pharmaceutical solutions by filtration although having advantages over other processes offers many serious objections. The list of thermolabile substances is increasing and dispensing them in sterile solutions with maximum activity is a difficult one. A faulty autoclave may be more easily detected than a faulty filter, a slight leak in a filter can only be detected by a bacteriological test of the filtrate. Most pharmacopæias which admit the process of filtration for sterile products include a sterility test of the final product. The major objections to the use of the bacterial filter are: (1) Mechanical construction defects—weakness of joints of connecting parts. (2) Adsorption—lowering of the concentration of the filtrate by adsorption of the medicament on the filtering ma-This varies with the medicament present. Adsorption will generally be at a minimum with collodion membrane filters and at a maximum with kieselguhr filters. (3) Yielding of alkali to the filtrate and alteration of pH of the latter—this may cause inactivation of certain substances as pituitary or insulin or may cause precipitation of alkaloids from solutions of alkaloidal salts. The Seitz filter may be open to this objection. (4) The vielding of small solid particles or floccules to the filtrate—the Seitz filter is open to this objection. It is often necessary to include a second clarifying filter in the system. (5) Difficulty of cleaning the filter-porous porcelain and kieselguhr filters are open to this objection. If it is necessary to heat the filter to a high temperature, cracking may result.

The following important factors must be considered in the mechanism of filtration: (1) The question of the actual size of the pores of the filter. By reducing the pore size filters can be made to

retain the bacteria but the rate of flow will be low, by increasing the thickness of the filter plate the pore size can be increased and consequently a greater rate of flow. The latter type generally has the disadvantage of high adsorptive properties. (2) The factor of surface equilibria between the substance of the filter and the dispersed phase (the bacteria) owing to adsorption and cohesion. (3) The reaction between the filter material and the continuous phase of the solution, in some cases causing swelling of the filter material. The Seitz filter is said to owe its efficiency to the latter principle.

The new 5/3 sintered or frittered glass filter seems to possess advantages over the older types of bacterial filters such as the Pasteur-Chamberlain, Berkfeld, Doulton and Mandler. Although the author cites no experiments regarding its power of adsorption, he states the apparent advantages of the Jena 5/3 filter to be: (a) it is made entirely of high resistant glass, the filter being fused into the glass holder, which reduces one risk of leakage. (b) It can be treated as glass, perfect cleaning being accomplished by using a concentrated solution of sulfuric acid heated to 80 degrees C. and containing a little sodium nitrate. Neither the efficiency or pore size is affected by this treatment.

In making the estimations of the size of the maximum pores for comparison with the maximum pore size of the older type filters the method of Bechold was used. This consists of pouring pure ether into the filter and forcing air through the plate from below; by placing a mercury manometer in the circuit it is possible to note the pressure at which the first bubbles of air appear through the plate. Bechold's formula for determining the pore size and a drawing of the apparatus are included in the article.

Using this method the Jena 5/3 filters compared favorably in maximum pore size with the other filters tested, more consistent results were obtained however by using carbon tetrachloride than with other. Tabulations of the pore sizes of the Pasteur-Chamberlain, German Berkfeld, Mandler, British Berkfeld, Doulton, Berlin Porcelain Co. (Massen type) and the Jena 5/3 type filters are included. From these tabulations the pore values of the L 5 Pasteur-Chamberlain and the porous porcelain English Doulton filters appear approximately equal—about 1.3 μ , while the Mandler (fifteen pound) has a value about 2.6 μ . The Jena 5/3 shows pore values between 3.2 μ and 0.74 μ .

tl

O

g

tr

d

of

cl

From bacteriological tests the Jena 5/3 filters were found efficient when the pore value did not exceed 2.5 μ .

In making the bacteriological tests, nine Jena 5/3 filters were assembled, autoclaved at 115 degrees C. for one hour, 100 mil quantities of water containing Chr. prodigiosum were drawn through and the filtrate tested for sterility. This organism was chosen because of its small size and ease of cultivation. The apparatus was next rinsed with water, again autoclaved and the filtration repeated. Each filter was used and autoclaved fifty times.

From the results of the sterility tests as shown in the table it was found that those filters which stopped the organism on the first filtration did so on every subsequent occasion, autoclaving fifty times did not affect their efficiency. Moreover the filters, although cleaned five times with the strong sulfuric acid solution, at the end of the operation showed no change in maximum pore size. In these tests several of the older types of filters were broken while the nine Jena 5/3 filters were still intact, showing sufficiently strong construction to withstand autoclaving and general handling.

H. P. F.

The Preparation of Dextrose Solution for Intravenous Administration. W. J. Elser and R. G. Stillman, Am. J. Clin. Path. 7, 221, 307 (1937). An investigation into the causes of reactions following the intravenous injection of dextrose solutions reveals the following factors: water not freshly distilled; particulate matter found in the solution; improperly prepared rubber tubing; too rapid or irregular injection of the fluid; unskillful performance of the operation; susceptibility of the patient; and a too high or a too low temperature of the solution. The following factors, though said to be the cause of the reactions, are probably of little importance: the pH and the age of the solution and the use of glass other than Pyrex. A method is given for the preparation of a 50 per cent. solution of U. S. P. dextrose which gives a solution that is clear, colorless and sterile and does not produce reactions. The features of this method are the use of only freshly distilled water, produced in a properly designed and operated still (double or triple distillation is not essential), and the clarification and sterilization of the solution in this method is by filtration through a bacteria-proof filter (Berkefeld). The dextrose

is at no time subjected to the action of heat which brings about changes in this material. This 50 per cent. solution has been shown to possess bactericidal power against the vegetative forms of the common pathogens.

L. G.

The Sterilization of Aqueous Zinc Sulfate and Potassium Aluminum Sulfate Solutions. A. Torstensen-Dalsgaard, Farm Tids 271 (1936), through Pharm. Zentralhalle 78, 646 (1937). Aqueous solutions of zinc sulfate and alum produce precipitates when heated for more than five minutes at 80 degrees C. in ordinary glass. The zinc salt can be autoclaved even for two hours in Pyrex or Jena glass without alterations of the pH. Alum solutions (10 per cent.) in Pyrex or Jena glass produce a precipitate when heated at 100 degrees C. or when autoclaved but not when heated for five minutes at 80 degrees C. Eye drops containing alum may be made by sterilizing concentrated solutions for five minutes at 80 degrees C. and diluting with autoclaved water.

Giemsa Preparation for Staining Blood Films. R. Hewitt, Science 86, 548 (1937). There have been many varieties of blood stains proposed, and each stain has its peculiar advantages. In a large proportion of clinical laboratories, where many blood films are stained daily, Wright's stain is generally found to be satisfactory and is relatively inexpensive. However, some laboratory workers prefer the more precise results obtained from other stains in the Romanowsky series. Giemsa's modification has been found to be very satisfactory as a routine blood stain as well as for blood parasites. Its chief disadvantage is its cost.

For several years the author has prepared blood films containing avian malarial parasites, using the Giemsa method exclusively. Ready prepared solutions of Giemsa generally cost about \$12 per eightounce bottle. The cost of eight ounces of prepared Wright's stain, on the other hand, is but \$2.85. This difference in price means a great deal to private laboratories and to many institutions. Some workers, even though preferring the Giemsa method, use Wright's stain regularly to reduce the expense involved.

of

R

B

The following method for preparing Giemsa stain, although not conforming to the usual technique suggested, has been found to be very satisfactory by the writer. The resulting stain costs about \$3.25 per eight ounces and gives uniformly well-stained blood films. In laboratories where a great deal of stain is used the cost of Giemsa so prepared is little more than Wright's.

Azur II—Eosin	3.0 gms.				
Azur II	o.8 gms.				
Glycerin (c. p.)	250.0 gms.				
Methyl alcohol	250.0 gms.				
Absolute (neutral), acetone free.					

Dissolve the Azur II and Azur II-Eosin in the methyl alcohol in a Erlenmeyer flask. Shake well for fifteen minutes, add glycerin, shake for ten minutes and filter through a moderately fine grade of filter paper. Collect the filtered stain in a bottle and discard the undissolved residue.

There is generally quite a bit of stain that does not dissolve. This, however, seems to make very little difference in the character of the resultant stained blood films. Results have been found equally satisfactory with human blood and avian blood. Malarial parasites are brought out sharply with distinct differentiation of chromatin and cytoplasm.

L. G.

Pneumonia Due to Type V Pneumococcus. M. B. Rosenbluth and M. Block, Arch. Internal Med. 60, 567 (1937). Of 1850 cases of pneumococcic pneumonia observed during the past eight years, 3.5 per cent. were type V. The mortality rate in thirty-nine cases of serum-treated type V pneumococcus pneumonia was 26 per cent. compared with 29 per cent. type V cases not treated with serum.

L. G.

Community Provision for the Serum Treatment of Pneumococcic Pneumonia. Report of the Committee on Public Health Relations of the New York Academy of Medicine by a Special Sub-committee. R. L. Cecil, J. G. M. Bullowa, H. T. Chickering and E. H. L. Crowin, Bull. N. Y. Acad. Med. 13, 557 (1937). This report covers pneu-

monia statistics, the history of pneumonia research, organization of pneumonia control in Massachusetts and New York, serum production in New York City and recommendations for further steps by the community to control the disease. The variation in the prevalence of pneumococcus types from year to year makes it difficult for health laboratories always to have adequate amounts of the different serums. Type I is consistently most prevalent. Second place has been held at various times by types III, VII and VIII.

L. G.

Gas Gangrene—Its Prevention and Treatment. B. Malone, J. Tennessee M. A. 30, 402 (1937). The polyvalent sera marketed in this country are not reliable. They were found to be unreliable in the treatment of gas gangrene.

Diphtheria Immunization: New Diluted Shick Fluid. C. R. Merrilless, M. J. Australia 2, 251 (1937), through J. Am. Med. Assoc. 109, 1492 (1937). Five thousand children were tested with a new diluted Shick toxin injected intradermally in the usual way in doses of 0.2 cc. The diluent consists of 8.4 gm. boric acid, 15.9 gm. Na Cl, 5.7 gm. borax and 5 gm. purified gelatin in two liters distilled water. A normal Shick reaction with the new preparation presents on the fifth day a striking patch of redness on a normal skin with practically no infiltration, no areolar paleness, and no tenderness or pain.

Accidental Smallpox Vaccination and Eczema Vaccination. G. W. Graves and C. Dowman, New York State J. Med. 37, 1833 (1937). Two cases of eczema vaccinatum, one of which was fatal, were reported. The presence of eczema in the individual or in members of the family is recommended as a contraindication for vaccination.

Diphtheria of the Skin. I. Pincus, New York State J. Med. 37, 1938 (1937). A case of diphtheria of the skin (a rare condition) was finally diagnosed after eight weeks. Administration of diphtheria antitoxin and application of 1:5000 potassium permanganate solution resulted in an uneventful recovery.

L. G.

Antiseptic Properties of Human Milk. H. Dold, E. Wizemann and C. Kleiner, Z. Hyg. Infektionskrankh 119, 525 (May, 1937), through J. Am. Med. Assoc. 109, 1640 (1937). Human or cow milk added to an equal volume of agar did not support the growth or allowed only slight growth of B. diphtheriae, Staph. aureus, B. coli, B. prodigiosus, B. pyocyaneus, B. anthracis, streptococci, and unidentified wild yeast. The factors in human milk inhibiting bacterial growth ("inhibins") were inactivated by heating at 56 degrees C. for thirty minutes or by standing twelve to twenty-four days at 5 degrees C., but not by repeated freezing and thawing. The "inhibins" in cow's milk were not inactivated by heating at 80 degrees C. for seven minutes but were destroyed by heating at 85 degrees C. for seven minutes. Attempts have not been made to identify the natural antiseptics.

The Choice of Disinfectant for Use in Hospitals. R. Hanne, Pharm. Ztg. 82, 957, 975 (1937). Disinfectants for hospital use are considered. In their purchase, factors to be noted are: toxicity, potency, cost, odor, color, effects on skin and instruments, stability, and the fact that the disinfectant action may be inhibited when in use.

L. G.

Misce et Fiat Collyrium. R. M. Nacca, Druggists Circular 81, 18 (1937). The preparation of solutions for use in ophthalmology are considered, emphasis being placed on isotonic eye solutions. The amount of salt necessary for rendering solutions isotonic is given. Two methods are described for buffering eye solutions. Formulae are presented for some of the more commonly used medicinals in ophthalmology.

L. G.

Sulfanilamide and Another Group of Diseases. Science Supplement 86, 12 (1937). Reports of successful use of sulfanilamide in a new group of diseases were made by Drs. Perrin H. Long and Eleanor A. Bliss, of the Johns Hopkins Hospital, Baltimore, at the meeting of the Southern Medical Association in New Orleans. The

first American report of the use of this powerful remedy was given by these same investigators at the meeting a year ago. At that time they reported cures of erysipelas, childbed fever and other strepto-coccus-caused ailments with sulfanilamide treatment. This year they reported its healing effect in the painful and generally stubborn illness due to infection of the urinary tract. Cases that had lasted for from five to seven years and in which all other forms of treatment had failed were cured by sulfanilamide. The drug, Dr. Long said, kills practically all of the germs which commonly infect the urinary tract.

The all important factor in treating these conditions is to obtain the proper concentration of the drug, that is, to get enough of it into the urinary tract to kill the germs. The amount necessary can be determined by laboratory tests. Laboratory tests, also, can show whether or not sulfanilamide will cure cases of streptococcus infections, the condition for which the drug was first tried. There are many varieties of streptococci. That not all are susceptible to the action of the drug is shown by a research reported by Drs. Long and Bliss and their associate, W. H. Feinstone.

Identification of the particular streptococcus that is causing illness in a particular patient is of prime importance. The investigators described how these germs may be identified and which one can be killed by sulfanilamide. Group A and Group B hemolytic streptococcal infections in man can be cured by the drug. Good results have been obtained in treating Group C hemolytic streptococcal infections in dogs and horses. Group D hemolytic streptococcal infections in man are unaffected by the drug. Alpha streptococcal infections, the kind found in the heart disease known as subacute bacterial endocarditis, are in general not affected by sulfanilamide treatment.

Cases of gonorrhea cured within three days by sulfanilamide were reported by Professor E. P. Alyea, of the Duke University School of Medicine, at the meeting. Professor Alyea treated 158 patients with this drug and four-fifths of them made rapid recoveries. Out of 1000 cases treated by the generally accepted method of a few years ago, nearly half had complications. Sulfanilamide was only introduced into American medicine a year ago and so far 830 cases of gonorrhea treated with it have been reported in scientific journals. In these 830 cases not a single complication of gonorrhea has been reported. Professor Alyea called this "a most striking result."

SOLID EXTRACTS

By Ivor Griffith, Ph. M., Sc. D.

Despite the form in which this information is presented it may be accepted as trustworthy and up-to-date. Original sources are not listed but they may be obtained upon request.

Garlic, as a medicinal, has enjoyed throughout the ages a considerable reputation. It is a sedative and alterative. But its real service to that portion of the human race that appreciates it, is as a savor, not as a drug or a scent. Five pounds of garlic are used daily in the Waldorf Hotel kitchens. Every week, Alfredo Maspero, the garlic king, handles for New York about twenty-five tons of the bulbs. He *knows* by rich experience, that garlic belongs to the lily family. Most of America's garlic (twenty million pounds a year) is grown in California.

Farmers hate garlic, because its odor contaminates milk, even though the cow does nothing worse than smell the weed.

Singers love it because it helps the voice, hence the success of cantors and Italian tenors in the field of music!

Actually, garlic, discriminately used, is the chef's best friend and the epicure's fond delight.

And today, since a gurgle and a gargle, with Dakin's solution, hides its announcing malodor, the garlic gourmet may safely ride the trolley without a grain of guilt.

Thymol has considerable application in medical practice, as an antiseptic, alone, or tied to iodine, as thymol iodide. Latterly, however, it is finding a wider range of usefulness in the industries. It is of value in preventing mildew growth, having been recommended for this purpose some time ago, by the Scientific Section of the National Paint, Varnish and Lacquer Association, especially for indoor and

industrial paints where the use of highly toxic metallic salts would not only be undesirable, but dangerous.

The paper industry has also found thymol of value as an antimold agent and its use seems to be increasing in that industry. It also has some stabilizing value such as in soaps and in face creams which may have a tendency to discolor and become rancid.

London has finally completed the terminal chloramination of its entire river-derived water supply, totalling about 250 million gallons a day. In good English, it might be now said with respect to the London supply, that both autochthonous and allochthonous pollutions have been mastered by physical and chemical antidotes.

At the present time, for days on end, every gallon of Aqua Londoniensis is 100 per cent. pure on a *coli* basis.

Which is quite an achievement! (Of course, we must recall that, despite the traditional tie-up of the Britisher to his bath, the average per capita consumption of water by the American city dweller is much higher than for the average Londoner.)

Many persons believe that typhoid-free water, containing other bacteria such as the b.coli is not a health hazard. They point out the fact that our intestinal tract abounds with such bugs.

But think again! Those waters with a high colonic count may be used to dilute and cool a young baby's bottle or the beef tea of an invalid. Should such be not consumed at once the poor person does not pay for replacement, but warms up the mixture, with the poisons resulting from the growth of these organisms, and possibly teeming with bacterial life, in other words, a laboratory culture. This is something to pause and ponder over, that a single organism may become two in twenty minutes or a quarter of a million in six hours, and that many of the filtrates derived from substrates of common organisms are definitely poisonous and dangerous to man.

Should a prescription for USNEA be brought to the modern pharmacy, there might be some difficulty in securing the drug. This is how Pomet's History of Drugs (early eighteenth century) describes it:

Casually we are told that supplies of it are usually obtained from the public executioner. The skull, hair and urine, and other parts and secretions are also recommended for various purposes, but the most interesting product from this source is our usnea, consisting of the moss which grew upon the human skull, particularly from individuals who had died violent deaths. If you wonder where supplies of such a drug could come from the question is answered by recalling that in England for centuries the bodies of criminals who were hanged were suspended in chains at the crossroads as a warning or deterrent to criminals. The text states casually that "The English Druggists bring these heads from Ireland, where they frequently let the bodies of criminals hang upon the gibbet until they fall to pieces, and you may see in the druggists' shops of London some of the heads entirely covered with moss."

Germany is undoubtedly the greatest nation in Europe on applied research. The nature of the people helps make this possible. It took fifteen years of hard, tedious plodding and 30,000,000 gold marks to produce the formula for aniline dyes.

The mammoth German dye trust is the I. G. Farben Industrie Co. at Ludwigshafen. Here buna and synthetic gasoline, products of coal, were discovered and developed. Germany being a country with plenty of coal, but not very good climate or soil, is fortunate that her scientists have been successful in working with coal for so much of her synthesis.

Here they have developed ceramic ware, similar to beetle ware, but cheaper and more transparent; molybdatrot, a new colored paint as fast as lead chromide and not subject to darkening.

They developed I. G. Wax, the use of which is now broadening so rapidly. It is used to give a better surface to paper; the quality of pulp or rag going into the paper can be of a considerably lower grade if I. G. Wax is mixed with it. This product also develops more beautiful colorings when mixed with paraffin and is the substance that prevents paraffin candles from bending in warm weather.

Another new product developed is locron, used to make cloth and building materials fireproof. A method has been developed to add pigment to it so that it can be applied as paint to furniture and woodwork—painting and fireproofing in one operation. At present the cost is twice that of ordinary paint, but it will decline with production.

E

C

C

C

D

Fu

Ge

The Kaiser Wilhelm Institute for Wood Research at Ebers-walde is doing considerable work with plywood impregnated with metals, and has developed a beech plywood impregnated with bakelite which is very strong. They have also developed what they call a radium wood which has been impregnated with oil. They have developed a new wood for beer barrels. This is a plywood, the center of which is beech, with both surfaces oak. The reason for the development of this wood was to enable them to save oak, but, through this plywood, they have also obtained a barrel that wears better.

Here they use the roots of pine trees, which are being dug up all over Germany, to obtain resins. They also use pine for turpentine and artificial silk. There is a new wood development in flooring, perfected in the United States, whereby the floor and the carpet are laid in one piece, joined together like a plywood. No dust can come between, and, inasmuch as dust causes much of the wear in carpets, this point was considerably stressed. It costs less to lay than when each material is handled separately, and it has other advantages which they believe will ultimately put it in general use for homes, office buildings and auditoriums.

INDEX TO VOLUME 109 OF THE AMERICAN JOURNAL OF PHARMACY

AUTHORS, 1937

(A)-	-Abstract	
(B)-	-Book Rev	riew
(E)-	-Editorial	

PAGE	PAGE
Batt, W. G.— Variations in the Electrophores- tic Behavior of Gelatin Pro- tected Silver Halide Sols 29	Greenbaum, Frederick R.— New Method for Iodation and Chlorination of Vegetable and Animal Oils
Calvert, Ralph— Pharmacal Arithmetic (B) 161	New Water Soluble Theophyl- line Compound 550
Chenoweth, Mary North— The Curious Case of Sir John Hill	Greene, L. Wilson— Chemical Microscopy of Fats and Waxes
Cohen, Isadore— An Outline of General Physiology (B)	Griffith, Ivor— Arsenomania (E) 65
Dunn, Marin S.— Appearance of N. F. VI Crude Vegetable Drugs Under Ultra-	Digested—But Hard to Swal- low (E)
violet Light	Gullibles' Travails (E) 454 Hush (E) 62 In 1936 (E) 6
Notes on the Effects of Certain Glandular Products Upon Plant Growth	In Praise of Good Molars (E) 457 I Went in Search of Flowers
Fulton, Charles C.— The Principal Chemical Tests for Morphine	
Gershenfeld, Louis— Prontosil and Prontylin (A) 47 Recent Developments in the Field of Antisepsis (A) 92	272, 336, 389, 449, 485, 531,
Recent Reports on Penumonia and the Pneumococcus (A) 432	Texts From the Textile Trades 18
Recent Reports on Prontosil, Sulfanilamide, and Related	This Cosmetic Democracy (E) 340 What Is Truth? (E) 536
Compounds (B) 367 The Wassermann, Kahn and	Words, Words, Words (E) 586
Other Serodiagnostic Tests for Syphilis (A)	
Goodrich, Forest J.— A Study of Struthiopteris Spi-	Disease? 348 Some Little Bug Is Going to
cant	Get You! 502

PAGE	PAGE
Harter, Elmer E., Jr.— New Food Legislation in Penn- sylvania	Prout, William A.— An Historical Sketch of the House of Solomons in Savan-
Heilmeyer, Ludwig— The Treatment of Iron-respon- sive Anemia With Iron Ascor-	Ramanuskas, Peter P.—
binate, and Contributions to the Mechanism of the Effect of Iron and to the Iron De- ficiency Disease Problem (A) 419	Cosmetic Cleansing Creams 131 Simon, Elsie— Experimental Determination of of the Amount of Benzedrine
Kelly, J. W.— The Detection of Ephedra Al-	in a Therapeutic Dose From Benzedrine Inhaler 343
kaloids in Small Quantities of Ephreda Kimmer, Wilton H.— Appearance of N. F. VI Crude	Simpson, Nathan A.— Experimental Determination of the Amount of Benzedrine in a Therapeutic Dose From
Vegetable Drugs Under Ul- tra-violet Light 498	Benzedrine Inhaler 343 Tice, L. F.—
Koff, Arnold— The Chemistry of Life (A) 86	Physical Chemistry (B) 160 Quantitative Pharmaceutical
Koozin, Estelle— A Study of Struthiopteris Spi-	Chemistry (B) 213 Ulan, M. S.—
Kramer, John E.—	Manual of Biological Assay- ing (B) 160
Principles of Pharmacy (B) 214 The March of Thyme 135	Van Itallie, Dr. L.— Outlook for an International
LaWall, Charles H.— Canning, Preserving and Pick- ling (B)	Pharmacopœia (A) 555 Viehoever, Arno—
Fads and Frauds in Foods and	Field Tests for Marihuana 589 Insect Control (A) 57
Drugs	Lipoids (A)
Iran and Iraq (B) 581 MacLaughlin, E. H.— A Practical and Inexpensive Vivarium for the Biology	Pharmacopœia (A) 555 Plant Hormones (A) 52 Progress in the Chemistry of
Laboratory 512	Life (A) 50
McArthur, C. S.— Concerning the Nature of Vitamin E	Textbook of Pharmacology (B) 276 The Chemistry of Life (A) 86 The Development of Daphnia Magna for the Evaluation of
McFadden, G. H.— Rhus Glabra	Active Substances 360 Translation of "Deutsche Archiv für Klinische Medizin,"
McMurray, R. L.— Rhus Glabra	Vol. 179, Pages 216-231 (1937), (A)
Milos, Charles— Detection of Cocaine in Mixtures of Cocaine and Nupercaine (1937)	Tschirch's Handbook of Pharmacognosy (B)
Parran, Thomas— The Aims of the Public Health	The Cosmetic Formulary (B) 277 Watson, E. M.— Concerning the Nature of Vita-

TITLE AND SUBJECT INDEX FOR 1937

(A)—Abstract (B)—Book Review (E)—Editorial (S)—Solid Abstract

PAGE	PAGE
Abiotoxine (S) 580 Acetone (S) 579	Anemia, The Treatment of Iron- responsive—With Iron Ascor-
Acetone, an Improved Method for Determining Ethyl Alcohol in the Presence of (A) 520	binate, and Contributions to the Mechanism of the Effect of Iron and to the Iron Deficiency Dis- ease Problem (A)
Acidosis (S) 391	Anesthetic Ether (A) 148
Adrenalin for Malaria (S) 450	Anesthetics (A)
Adrenalin, Stabilization of-Solu-	Aniline Dyes (S)
tions (A) 604	An Outline of General Physiol-
Adrenals for Adaptation, The Significance of the (A) 243	ogy (B) 583
Adsorption of Ions and the Physical Character of Precipitates	Anti-Blood Clotting Agent Clue to Thrombosis Treatment (A) 515
(A) 82	Antibody, The Structure of the
Aerogel (S) 158	(A) 197
Agar, Emulsion of Liquid Paraf-	Antigens (A) 84
fin With (A) 39	Antioxidant, Oat Flour as an (A) 78
Agricultural Insecticides (A) 245	Anti-rust Agents (S) 158
Alcohol Content of Extractive Preparations (A) 477	Antisepsis, Recent Developments in the Field of (A) 92
Allantoin (S) 155	Antiseptic Value and Pharmacol- ogy of an Improved Calomel
Allyl Aldehyde (S) 212	Ointment (A) 459
Aluminum Sulfate (A) 610	Antiseptics: A Comparative Study
American Association for the Advancement of Science and Asso-	of Laboratory and Practical
ciated Societies, Some Papers	Tests (A)
Read at the Atlantic City Meet-	Antiseptics, Longevity of (S) 579
ing of the (A) 84	Antiseptics in Obstetrical Practice (A)
American Larch, The Bark of (A) 248	Antiseptics, Relative in Vitro Ac-
American Mistletoe (A) 190	tivity of Certain—in Aqueous
Amidopyrine and Aspirin (A) 571	Solution (A) 154
Aminophyllin 550	Antiseptics, Scientific Advances
Aminophylline (A) 568	in 1936 in Disinfectants and 123
Ammonium Carbonate, Incompat- ibility Between Syrup of Tolu	Antiseptics and Wound Healing (A)
and—(A) 186	Ants and Their Control in Cali-
Ammonium Mandelate, Syrup of	fornia (A) 526
(A) 80	Argentine Ant Syrup (A) 529
Androstanedione, Preparation of —From Allo-Pregnandiol (A) 242	Antuitrin Growth II Apothecary Benham (S) 449

PAG		AGE
Apple Acid (S) 10	Benzedrine Inhaler, Experimental	
Aqua Londoniensis (S) 61		
Aqua Philadelphica (E) 11	Benzedrine in a Therapeutic Dose From	242
Arsenic (S) 6		343
Arsenic Poisoning, Fixing the	C (A)	322
Time of the Administration of	Benzyl Methyl Carbinamine S	0
Poison in a Case of Chronic (A) 19	3 K F	343
Arsenic, Report on (A) 31	Berries, The Food Value of (A)	202
Arsenomania (E) 6	Beryllium	58
Artemisias, Welsh and English,	Beverages Public Health Aspects	50
	of the Treatment of Water and	
Ascorbic Acid, A Color Reaction	-With Silver (A)	141
of (A) 44	Bezoar Stones (S)	212
Ascorbic Acid, Autoxidation of	Rile Pigments (S)	
(,	Biological Test Animal, Daphnia	
Ascorbic Acid, Oxidation of (A) 18	(A)	95
Aspirin and Amidopyrine (A) 57	Diology for Students of Fliat-	
Astrology (S) 20	macy (D)	582
Atropine and Syntropan, A Com-	Bismuth Salts, Color Changes in	
parative Study of (A) 38		
Audubon (S) 44		442
Auxine (S)		
Azobenzenesulfonamides, The Io-	Blond Hair (S)	101
dination of p-Aminobenzenesul- fonamide and Some Symmetrical 41	Blood Films, Giemsa Preparation	
	2	010
Bacteria (S)		
Bacterial Flora, The Relationship		
Between Oral and Gastric (A) 25	Blueberries (A)	
Bacterial Mounts, The Use of	body remperature (b)	-
Synthetic Resins in the Prepara-	Book Reviews 276,	_
tion of Permanent (A) 38		
Bacteriophage and Its Relation to	Bread Freezing (A)	
Oral Infection 13	Brewing Industry (S)	532
Bacterium (S) 33	6 Brillat-Savarin (S)	532
Baking Industry (A) 14	6 British Herbal	460
Barbituric Acids, Containing the	Brushless Shaves, Tested Formu-	
2-Methylallyl Group (A) 48	las (A)	
Barbiturate, Decomposition of-	Buckingham's Reagent	228
Solutions by Heat (A) 116	Bug, Some Little-Is Going to	
Bath Preparations (A) 18	5 Get You	
Beeswax, Chemical Constants of	Bunsen	58
(A) 79	Burns, Poisons Found in Blood	
Benzedrine, Experimental Deter-	Following Extensive (A) 2	255
mination of the Amount of—in	Burns, Tannic Acid-Silver Ni-	
a Therapeutic Dose From Ben-	trate Treatment of (A) 4	444
zedrine Inhaler 34	3 Buromin	25
Benzedrine, Hypertension Pro-	Cacao Hulls as a Source of Vita-	
duced With (A) 38	3 min D (A)	42

INDEX TO VOLUME 109

Calcium Ortho-Iodoxybenzoate in the Treatment of Arthritis (A) 448 Calculi (S) 211 Calex 25 Calgolac 25 Calgon 25 Calgonite 25 Calounel Ointment, Antiseptic Chlorine-Containing Compounds as Bactericidal Agents (A) Chlorine Therapy of Infectious Hemorrhagic Colitis (A) Chloropicrin (S) Chloropicrin (S) Cholesterol (S) Chrysopsis Graminifolia, Nutt (A) Cinchophen in Presence of Sodium	407 113 145 250 159
Use (A)	113 145 250 159
Calcium Gluconate Ampuls (A) 564 Calcium Gluconate Solutions (A) 478 Calcium Ortho-Iodoxybenzoate in the Treatment of Arthritis (A) 448 Calculi (S) 211 Calex 25 Calgolac 25 Calgon 25 Calgonite 25 Calounel Ointment, Antiseptic Method for Chlorine Cocktail (E) Chlorine-Containing Compounds as Bactericidal Agents (A) Chlorine Therapy of Infectious Hemorrhagic Colitis (A) Chloropicrin (S) Chloropicrin (S) Chloropicrin (S) Chloropicrin (S) Chloropicrin (S) Chloropicrin (S) Chlorophen in Presence of Sodium	113 145 250 159
Calcium Gluconate Solutions (A) 478 Calcium Ortho-Iodoxybenzoate in the Treatment of Arthritis (A) 448 Calculi (S) 211 Calex 25 Calgolac 25 Calgon 25 Calgonite 25 Calounel Ointment, Antiseptic Chlorine Cocktail (E) Chlorine-Containing Compounds as Bactericidal Agents (A) Chlorine Therapy of Infectious Hemorrhagic Colitis (A) Chloropicrin (S) Chloropicrin (S) Cholesterol (S) Chrysopsis Graminifolia, Nutt (A) Cinchophen in Presence of Sodium	113 145 250 159
Calcium Ortho-Iodoxybenzoate in the Treatment of Arthritis (A) 448 Calculi (S) 211 Calex 25 Calgolac 25 Calgon 25 Calgonite 25 Calounel Ointment, Antiseptic Chlorine-Containing Compounds as Bactericidal Agents (A) Chlorine Therapy of Infectious Hemorrhagic Colitis (A) Chloropicrin (S) Chloropicrin (S) Cholesterol (S) Chrysopsis Graminifolia, Nutt (A) Cinchophen in Presence of Sodium	145 250 159
Calculi (S)	250 159
Calex	159
Calgolac	159
Calgon 25 Cholesterol (S) Chrysopsis Graminifolia, Nutt (A) Calomel Ointment, Antiseptic Cinchophen in Presence of Sodium	211
Calgonite	
Calomel Ointment, Antiseptic Cinchophen in Presence of Sodium	
	,
Value and Pharmacology of an Bicarbonate (A)	184
Improved (A) 459 Cinnamon Industry in Ceylon (A)	
Camouflage (S) 452 Cleasing Creams	131
Cancer 128 Cobalt in Metallic Nickel, On the	
Cannabinol	147
Cummour Duracema (E)	14/
Canning, Preserving and Pick- ling (B)	332
Carbolic Acid Bands for Ants (A) 527 Cocaine and Nupercaine, Detec-	
Casein, Determination of, by For-	116
Titudion After Precipita Cocaine, Detection of-in Mix-	
tion With Acid (A) 83 tures of Cocaine and Nupercaine	
Castor Oil 67 Cod Liver Oil (S)	579
Castor Oil (S) 579 Cod Liver Oil, Vitamin A Con-	
Castor Oil Hydrogenated as an	
Ointment Base (A) 74 Coffee (A)	41
Ceramic Ware (S)	241
Ceresin	250
Cevitamic Acid, Autoxidation of	
-and its immortion by Sui-	-/-
phur (A)	131
in (A) 199 Colloids (S)	
Chediak's Dry Blood Reaction (A) 474 Collyrium (A)	
Chemicals or Germs Cause Dis- Color (S)	
ease? 348 Colors (S)	
Chemical Tests for Morphine, Color Changes in Ointments (A)	142
The Principal 219 Compounding The Art of (B)	
Chemistry of Life (A) 80 Contamination of Ice Cream (A)	
Chemistry of Life, Progress in the (1937), (A) 50 Consumer's Research (S)	
(193/), (11)	
Chemistry of Nutrition, The Pres- Copper Sulfate, Effect of (A)	
Chemistry of Nutrition, The Present Position of the (A) 263 Copper Sulfate, Effect of (A)	205
Chemistry of Nutrition, The Pres- Copper Sulfate, Effect of (A)	205

PAG	E	AGE
Cosmetic Cleansing Creams 13	Diphtheria Toxin (A)	522
Cosmetic Democracy (E) 34		
Cosmetic Dermatology (A) 48		
Cosmetic Formulary (B) 27		
Coumarin in Vanilla Extract (A) 52		348
Cranberries (A) 20		
Cricket (S) 15		123
Crotonic Aldehyde (S) 21	Disinfectant for Use in Hospi-	
Crystals (S) 33		613
Culture Media. The Prevention of	(A)	600
Drying-Out in (A) 20	(A)	003
Curious Case of Sir John Hill 46	(A)	78
Cyclohexylamine (S) 15		
Cytolysins (A) 46		103
Czapek's Solution 18	G B min Gminitarion	80
Daguerre's Old Tintype Photo-	Drug Market Committee Report.	
graphs (S) 10	5 1937	464
Daphnia (S) 33	Drying-Out in Culture Media,	
Daphnia 59	Prevention of (A)	201
Daphnia, as a Biological Test	Dupanol	21
Animal (A) 9.	Duroquinine	548
Daphnia, Cardinal Points in Favor	Eagle Antigens (A)	373
of (A) 26	Eczema Vaccination and Small-	
Daphnia Magna		012
Daphnia Magna, The Develop- ment of—for the Evaluation of	Eighteen Twenty-One, Ever Since (E)	2
Active Substances 36		=6T
Deceresol		301
Dehydroascorbic Acid, Vegetal	atin Protected Silver Halide	
Reduction of (A) 49%	Sols	29
Dental Powder (1937), (S) 210		493
Deodorants, Tested Formulas (A) 32	Emulsification Power of Infusions	
Dextrose Solution, Preparation	of Medicinal Plants, Reasearch	
of (A)	Francisco of Times 1 December 1877:41	243
Diet and Nephritis (A) 56;	Agar (A)	39
Diethylene Glycol (E) 494	Enterocrinin (A)	
Diethylene Glycol, Toxicity and	T	212
Fate of When Administered to Animals (A) 70		
Digested—But Hard to Swallow	in Small Quantities of Ephedra	36
(E) 112	Ephedra, Cultivation of, in South	
Digestion (A) 86	Dakota (A)	45
Digitalis (A) 149	Ergosterol, Irradiated, Treatment	
Digitalis Tincture (A) 480	of Psoriasis (A)	601
Dinitrophenol (A) 186		24
Dioxan, Toxicology of (A) 44	Tel - A (A)	148
Diphtheria Immunization (A) 612		
Diphtheria of the Skin (A) 612		520

PA	GE PAGE
Ethylene Dichloride and Carbon Tetrachloride Mixture for Ants	Friday for Fish (S)
(A) 5	
Ethylene Glycol (E) 4	
Euphylline 5	
Euthanasia, Small Animal—Vet- erinary Surgeons' Recommenda- tions (A)	Fur Names (S)
	49 Gallstones (S)
Evaluation of Germicidal Sub-	Gallstones (S)
stances (A) 5	25
Ever Since the Quiet Quaker Days	Gardinoi, W. A
of 1821 (E)	Garlic (S)
Extra Pharmacopœia I	08 Garlic (S)
Extracts and Tinctures, Prepara-	Gas Gangrene (A) 612
tion of (A) 4 Fads and Frauds in Foods and	phorestic Benavior of-Pro-
Drugs 1	15 Commission Oil (A)
Family Herbal (1937) 4	German Dye Trust (S) 617
Fats, Chemical Microscopy of	07
Fatty Fish, Oat Flour Improves Keeping Quality of (A) 5	Germs or Chemicals Cause Dis-
Federation of International Pharmacy (A) 5	ease?
Fever Treatment (A) I	Blood Films (A) 010
Fiber Stain, A Useful—for the Paper Chemist (A) 2	Plant Growth 9
First Chemist (S)	
Flavoring Agents, The Compara- tive Efficiency of the Commonly	Goiter and the Use of Iodine in
Used (A) I	gi Its Prevention (A) 269
Floor Polish	28 Gold (S) 104
Flowers, I Went in Search of (E) 3	Gonorrhea, New Antitoxin Prom-
Fluckiger's Reagent 2	ises a Cure of (A) 254
Fluorescence (S)	Gonorrhea, New Chemical Cure
Fluorine, Chemical Removes (A) 5	10 10 10 10 10 10 10 10 10 10 10 10 10 1
Fluorine Compounds, Toxicity of	Gout (3)
(A) 5	Gray Hair (S) 337
THE . TO	Grease Baits for Ants (A) 529
Food (S) 5	Grey Powder (A) 142
Foods and Drugs, Fads and	Guilibles Travails (E) 454
Frauds in I	15 Hand Lotion, Milky (1937) 27
Food Legislation in Pennsylvania 5	
Food Regulations (S) 5	Helenalin, From Helenium Au-
Food, What Happens to-in Your	tumnale (A) 43
	88 Hepburn Bill 115
Formol Titration, Determination	Hill, Curious Case of Sir John 460
of Casein by-After Precipita-	Hinton Test for Syphilis (A) 473
	83 Hormone, New (A) 572

PAGE	PAGE
Hormones (Plant), (A) 52	-, and Contributions to the
Hormones, Plant Growth (A) 66	Mechanism of the Effect of Iron
Human Milk, Antiseptic Proper-	and to the Iron Deficiency Disease Problem (A) 419
ties of (A) 613	Iron, Determination of (A) 569
Human Pathogenic Fungus as a	Iron, Determination of (A) 509
Saprophyte (A) 584	Isaac Newton (S) 533 Ispaghula, Psyllium and Related
Hush! (E) 62	Seeds (A) 46
Hydrocyanic Acid, The Detection	Ivy Poison (S) 579
of Inhaled (A) 521	I Went in Search of Flowers (E) 394
Hydrosol of Silver Iodide 30	Jorissen's Test (A) 265
Hypochlorites, Comparison of the Germicidal Efficiency of — of	Kahn, Wassermann and Other
High and Low Alkalinity (A) 145	Serodiagnostic Tests for Syph-
Ice Cream, Contamination of (A) 602	ilis (A) 468
I. G. Wax (S) 618	Karaya Gum (A) 40
Immunity 592	Karl Friedrich Mohr (S) 210
Immunization by Living and Path-	Kerosene and Pyrethrum Mixture
ogenic Staphylococci (A) 530	for Ants (A) 527
Immunology Practices and Tech-	Kidney Stones (S) 208
niques (A) 324	Kline Test for Syphilis (A) 472
Industrial Cancer (A) 198	Knop's Solution Modified 10
Infusions of Medicinal Plants, Re-	Labiatae, Development of the Mac-
search on the Emulsification	rogametophyte in Certain (A) 45
Power of (A) 243	Lactic Acid (S) 391
In 1936 (E) 6	Lactobacillus Acidophilus, The Role of—in the Treatment of
Inoculations, Preventive (A) 262	Disorders of the Digestive Tract
In Praise of Good Molars (E) 457	(A) 76
Insecticides (A) 56	(A)
Insecticides, Agricultural (A) 245	(A) 248
Internal Parasites, Control of (A) 204	Lead and Arsenic Insecticides (S) 275
International Pharmacopœia (A) 555	Lead Poisoning Rates Worst as
Intestinal Function (S) 485	a Hazard to Workers (A) 330
Iodation and Chlorination of Veg-	Lefort Test 225
etable and Animal Oils, New	Life Is a New Creation (E) 164
Method for 407	Life Is Colloid (E) 165
Iodine (S)	Lipoids (A) 54
roid Disease (A) 266	Liquid Paraffin, Emulsion of—
Iodine (S)	With Agar (A) 39
Iodine (1937), (A)	Liquor Antisepticus (A) 479
	Lithium Salts (S)
Iodized Salt (A) 271	Liver in Therapy (1937), (S) 485
Iodometric Determination of Sugar	Locron (S) 618
(A)	Lubricating Creams (A) 139
Ions, Adsorption of—and the Physical Character of the Precipi-	Luminal Injection Solutions (A) 381
tates (A) 82	Lycopodium Granuloma (A) 570
Iron Ascorbinate The Treatment	Marihuana 589
of Iron-responsive Anemia with	Molars (E) 457

PAG	E PAGE
Molybdatrot (S) 61	TAGE
Mild Tincture of Iodine (S) 53	
Mercuro-chrome (S) 534	Standardization of Tablets for
Micro-analysis (E) 492	Determining 415
Metaphylline 550	Methylenedisulfonic Acid and Its
Mono-piperazine Theophylline 551	Derivatives, A Study of (A) 379
Milk, Unpasteurized (A) 564	Microscopy, Chemical — of Fats and Waxes
Measuring Devices (S) 578	Milk and Nutrition (A) 325
Macrogametophyte, Development	Milk, Standardization of Tablets
of-in Certain Labiatae (A) 45	for Determining Methylene Blue
Maggots (1937), (S) 155	Reduction in 415
Maggots The Therapeutic Active	Millon's Test (A) 265
Principle of	mississippi, Drug Concernig and
peutics of (A) 254	Cultivation (A) 80
Magnesium Phosphate, Tribasic	Modern Archemy (5) 430
(A) 137	Mohler's Test (A) 265
Magnesium Sulfate as an Insecti-	More for Your Money (B) 277
cide (A) 318	Morphine, The Principal Chemical Tests for
Mal-de-mer (S) 336	Mose's Anointing Oil (S) 90
Malic Acid (S) 102	Mouth, Yeasts in the Normal 167
Mandelic Acid—Bactericidal Ac-	Muscle Nuclei, The Isolation of
tion (A)	(A) 246
Urinary Infections (A) 44	Mushrooms (S) 390
Mannick's Method for the Deter-	Mutual Flocculation Phenomenon 33
mination of Opium (A) 382	Natural Immunity 592
Manual of Biological Assaying	Nephritis and Diet (A) 563
(B) 160	Nerve Impulses, Chemical Trans-
Marme's Reagent 222	mission of (A) 51
Marquis' Reagent 230	Nicholl's Test (A) 265
Mechanism of Strychnine 285	Nicotine (S) 531
Mecke's Reagent 229	Nicotine Sulfate for Ants (A) 527
Menhaden Fish Oil, Vitamin D	Nicotine Thiocyanate (A) 605
Content of (A) 148 Mercurial Antiseptics, Mechanism	Nitrosophenylhydroxylamine (A) 569
of Inactivation of—Serum, and	Nose Drops (S) 156
Its Implications Regarding the	Nose Sense (1937), (S) 274
Possibility of Intravenous Anti-	Numerology (1937), (S) 208
sepsis (A)	Nutrition, The Present Position of
Mercury Vapor (S) 105	the Chemistry of (A) 263
Merthiolate (A) 207	Oat Flour as an Antioxidant (A) 78
Metallic Nickel, On the Colori- metric Determination of Co-	Obstetrical Practice, Antiseptics
balt in (A) 447	(S)
Metaphen (A)	Obsterical Practice, Antiseptics
	in (A)
Methemoglobin (A) 322 Methenamine on Sterilization, De-	in (A) 207
composition of (A) 380	Odor Developed (S) 273
	(-)

PAGE	PAGE
Of Little and Less (E) 495 Ointments, Color Changes in (A) 442	Philadelphia College of Phar- macy and Science, One Hun-
One Hazardous Day (E) 216	dred and Fifteenth Annual Com-
Onion (S) 212	mencement
Opium, Mannick's Method for the	Philadelphia Water (E) 114
Determination of (A) 382	Phosgene (S)
Optical Lenses From Plastics (A) 141	Phosphatase Test (A) 565
Organic Chemicals, Trends in-	Physical Chemistry (B) 160
From Petroleum and Natural	Physicians (S)
Gas (A) 138	Picric Acid-Alum Spray, Experi-
O'Rourke, Constance (S) 449	ence With the—in the Preven- tion of Poliomyelitis in Ala-
Ozone (S) 531	bama (A) 206
Palaemonetes 305	Piperazine Hydrate 550
p-Aminobenzene Sulfonamide, The	Piperidine (A) 481
Effect of—on Pneumococci in Vitro (A) 144	Pituitary Gland (S) 531
p-Aminobenzenesulfonamide, The	Pituitary of Cattle, The Compar-
Iodination of-and Some Sym-	ative Pharmacognosy of the An-
metrical Azobenzenesulfona-	terior and Posterior Lobes (A) 192
mides 415 Parachlormetaxylenol as a Pre-	Plant Growth, Effects of Certain Glandular Products 9
servative (A) 137	Plant Hormones (A) 52
Parahydroxybenzoic Acid, The	Plastics, Optical Lenses From
Detection and Determination of	(A) 141
—and Its Derivatives With	Platinum (S) 104
Special Reference to Their Dis- tinction From Salicylic and Ben-	Plywood Impregnated With Met-
zoic Acids (A) 264	als (S) 618
Parthenocarpy Induced by Pollen	Pneumococcus, Recent Reports on
Extracts (A) 143	Pneumonia and the (A) 432
Pellagri's Test 234	Pneumonia
Pennsylvania, New Food Legisla-	
tion in 538	Pneumonia, Community Provision for the Serum Treatment of
Percolation, Theoretical and Prac-	(A) 611
tical Investigation on the Proc-	Pneumonia Deaths, Chemical
ess of (A)	Promises to Reduce (A) 325
Perilla Oil (S)	Pneumonia and the Pneumococ-
esses Used for (A) 198	cus, Recent Reports on (A) 432
Peroxide of Hydrogen 26	Pneumonia Serum Obtained From
Pertussis Endotoxin in the Treat-	Rabbits, Better (A) 254
ment of Whooping Cough (A) 554	Podagra (S) 209
Pharmacal Arithmetic (B) 161	Poisonous, Plants Made by Sele-
Pharmaceutical Education in China (A)	nium Absorbed From the Soil (A)
Pharmaceutical Recipe Book (B) 107	Poliomyelitis, Recent Studies on
Phenol, The Colorimetric Method	(A) 206
for the Determination of Traces	Polio, Zinc Sulfate Pontocain
of—in Water (A) 186	Solution Urged as a Preventive
Phenolphthalein Studies (A) 456	of (A) 518

PAGE	PAGE
Poisons Found in Blood Following Extensive Burns (A) 255	Psoriasis, Treatment of—With Massive Doses of Crystalline
Pollen Extracts, Parthenocarpy,	Vitamin D and Irradicated Ergosterol (A)601
Induced by (A) 143	Psyllium, Ispaghula and Related
Popular Science (E) 4	Seeds (A) 46
Population of China (S) 103 Potassium Chlorate in Dentifrices	Public Health Service, The Aims
(A) 321	of the 124
Potassium, Determination of—as	Purists (S) 389
Potassium Silver Cobaltinitrite	Pyrethrum (A) 56
(A) 342	Pyrethrum and Soap Solution for
Potassium Silver Cobaltinitrite,	Ants (A) 528
Determination of Potassium as	Quantitative Pharmaceutical Chem-
(A) 342	istry (B)
Potatoes, Vitamin C Content of	Quick and the Dead (E) 492
(A) 317 Prescription Room Equipment, A	Quinine Sulfate as a Fluorescent Indicator for Precipitation Re-
Check on (A) 84	actions (A) 318
Preserving Powder (S) 391	Quinine, Wars Won With-In-
Preserving Solution for Green	stead of Artillery (A) 327
Plants (A) 91	Rabbits, Better Pneumonia Serum
Prestone (E) 492	Obtained From (A) 254 Rabies Found to Occur at Any
Principles of Pharmacy (B) 214	Season of Year (A) 328
Procaine Solutions, Adjustment	Rabies Menace; Increases; Med-
of pH Value of (A) 379	ical Men Urge Action (A) 326
Prontosil (A)	Rain (S) 273
Prontosil and Prontylin (A) 47	Raisin Seed Oil (A) 517
Prontosil, Sulfanilamide and Re- lated Compounds, Recent Re-	Resins, Synthetic, The Use of-
ports on (B) 367	in the Preparation of Permanent Bacterial Mounts (A) 380
Prontylin and Prontosil (A) 47.	Rhaponticum and Rhubarb (A) 561
Prontylin, The Effect of p-Amino-	Rheum Rhaponticum (A) 561
benzene Sulfonamide on Pneu- mococci in Vitro (A) 144	Rhubarb and Rhaponticum (A) 561
Propadrine Hydochloride (A) 537	Rhus Glabra
Prophylaxis Against Pertussis	Rhus Glabra, Preliminary Chem-
and Chicken-pox, Passive (A) 207	ical Investigation of the Berries
Prophylaxis Against Poliomyeli-	of (A) 77
tis, Passive (1937), (A) 206	Roach Exterminator (1937) 25
Propylene Glycol (E) 492	Rose Water, Tests for Quality in
Propylene Glycol, Toxicity of (A) 478	the P. H. V. (A) 43
Protamine Zinc Insulin (A) 100	Rotenone and Pyrethrins (A) 56
Protein, Non-living—Causing Dis-	Rubber Caps for Vaccine Bottles (A) 323
ease in Animals (A) 147	Rush, Dr. Benjamin (S) 272
Protocatechuic Acid (S) 337	Sabouraud's Media
Provitamin and Vitamin A. Vital	Saccharomyces
Substances in Food and Medicine (1937), (A) 50	
	Salt (S)
Prudists (S) 392	Salt Diet (S) 389

PAGE	PAGE
Santonin in English and Welsh Artemisias (A) 38	Sodium Formaldehyde Sulfoxalate (A)
Sarracenia (E)	Sodium Mandelate as a Urinary
Sauer's Vaccine (A) 554	Antiseptic (A) 523
Scabicidal Drugs: An Experi-	Sodium Perborate (S) 451
mental Study 123	Sodium Perborate Therapy (A) 480
Scabies, Treatment of, With Ben-	Sodium Phenobarbitone (A) 606
zyl Benzoate (A) 481	Sodium Salicylate, Stabilization
Sciosophy (S) 209	of Solutions Containing-and
Sea-sickness (S) 336	Sodium Bicarbonate (A) 114
Selenium, Plants Made Poison-	Sodium Urate (S) 209
ous by—Absorbed From the Soil (A)	Solid Extracts58, 101, 155, 208, 272, 336, 389, 449, 485,
Serodiagnostic Tests for Syphilis	531, 575, 615
(A) 468	Solomons, House of 72
Serum (A) 207	Some Little Bug Is Going to Get
Serum Therapy (A) 75	You! 592
Serum Treatment of Pneumonia,	Snuff (S) 274
Community Provision for (A) 611	Spectograph (E) 495
Shick Fluid (A) 612	Spectrum (S) 533
Ships May Get Sick Through	Spray Materials: Recent Devel-
Germ Attacks (A) 196	opment in Fungicides (A) 46
Silver Halide Sols., Variations in	Stabilization of Solutions Containing Sodium Salicylate and
the Electrophorestic Behavior of Gelatin Protected 29	Sodium Bicarbonate (A) 114
Silver Picrate Treatment of Vag-	Stain, A Useful Fiber-for the
inal Trichomoniasis (A) 524	Paper Chemist (A) 247
Silver, Public Health Aspects of	Stanley's Discovery (A) 98
the Treatment of Water and	Staphylocci, Immunization by
Beverages With (A) 141	Living and Pathogenic (A) 530
Smallpox Vaccination and Eczema Vaccination (A) 612	Staphylococcus Aureus, Cultur-
Smelling Game (S) 577	ing (A)
Smell-shock (S) 577	Air and Steam and of Super-
Smog (S) 60	heated Steam (A) 529
Soap Preservatives (A) 244	Sterilizing Lamps (A) 191
Soap Wash for Ants (A) 527	Sterols (A) 242
*	Street Dust, Isolation of Tetanus
Social Security Plan (S) 60	Bacilli From (A) 530
Sodium Arsenite for Ants (A) 528	Struthiopteris Spicant, A Study of 412
Sodium Bicarbonate, Cinchophen	Strychnine (A) 569
in Presence of (A) 184	Strychnine, Mechanism of 285
Sodium Bicarbonate, Stabilization of Solutions Containing Sodium	Sugar, Iodometric Determination
Salicylate and (A) 114	of (A) 35
Sodium Chlorite as a Volumetric	Sugar Makes Alcohol Less Toxic
Oxidizing Agent (A) 81	(S) 338
Sodium Fluoride and Pyrethrum	Sulfanilamide (A) 613
for Ante (A)	Sulfanilamide Flixir (F) 402

TTTTTTT

DAGE	DAGE
Sulfanilamide, Prontosil, and Re-	Theophylline Iconropanolamine 552
lated Compounds, Recent Re-	Theopreparel
port on (B) 367	Theopropanol 553 Thrombosis Treatment, Anti-blood
Sulfate Solutions (A) 610	Clotting Agent Clue to 515
	Thyme, The March of 135
Sulfonamide Compounds, Chemo- therapy of Virus Diseases With	Thymol (S)
(A) 52I	Thyroid Disease, Iodine in the
Sulfur Bands for Ants (A) 527	Prevention of (A) 266
Sunburn (S) 392	
Sunburn Protection Measured (A) 94	
Synthetic Aliphatic Penetrants	Tincture of Digitalis (A) 480
(A) 606	Tinctures, Preparation of Extracts and (A)
Syntropan and Atropine, A Com-	Tintype (S)
parative Study of (A) 383	Titanium Dioxide, Dispersing (S) 580
Syphilis (E)	m m
Syphilis (S) 101	a-Tocopherol, The Thermal De-
Syphilis127	composition of (A) 366
Syphilis, Efficiency of State and Local Laboratories in the Per-	Toothpaste (A) 565
formance of Serodiagnostic	Tooth Powder (S)
Tests for (A) 259	Toxicity of Propylene Glycol (A) 478
Syphilis, Ide's New Color Test	Traffic in Health (B) 488
for (A) 259	Tri-Calcium Phosphate 25
Syphilis, Wassermann, Kahn and	Tri-Calcium Phosphate (A) 519
Other Seriodiagnostic Tests for	Trichomonas Vaginitis, New
(A) 468 Syrup of Ammonium Mandelate	Treatment for (A) 76
(A) 80	Trichophytosis, Copper Sulphate
Syrup of Tolu, Incompatibility	Treatment of (A) 588
Between—and Ammonium Car-	Triethanolamine 27
bonate (A) 186	Triethanolamine (A) 482
Tannic Acid-Silver Nitrate Treat-	Triethanolamine and Other Etha-
ment of Burns (A) 44	nolamines (B) 106
Tartar Emetic and Sugar Syrup	Triethanolamine, A Rapid Method
for Ants (A) 529	for the Determination of (A) 248
Tergitol	Triethanolamine Salts of Fatty Acids (A) 478
Against With Tetanus Toxoid	Trigamine
(A) 75	Tri-Sodium Phosphate 25
Tetanus Bacilli, Isolation From	Tschirch's Handbook of Phar-
Street Dust (A) 530	macognosy (B) 162
Tetanus Toxoid. Active Immuni-	Tuberculin Test (A) 251
zation Against Tetanus With	Tugwell Bill
(A)	Typhoid Fever, Vaccination
	Against 549
Tetraphosphoric Acid 5 Textbook of Pharmacology (B) 276	Typhoid-free Water (S) 616
Texts From Textile Trades 18	
Theophylline (A)	Typhus Fever (A) 530
Theophylline Compound 550	Ultra-violet Light 57
Theophylline Ethylenediamine 550	Ultra-violet Light, Appearance of N. F. Crude Vegetable Drugs. 498
incopily in the Langue in the control of the contro	14. 1. Citide Vegetable Diugs., 490

United States Dispensatory, The New (B) 109	Vitamin D, Treatment of Psoriasis (A) 601
Urinary Lithiasis, Vitamin A De-	Vitamin E 544
ficiency in the Eye Correlated	Vitamin E, Wheat Germ Oil as a
With (A) 604	Valuable Source 544
Urine (S) 103	Valuable Source
Useful Plants and Drugs of Iran	Vitamins (S) 337
and Iraq (B) 581	Vivarium 512
	Wadsworth's Antigen (A) 474
Usnea (S)	Wagner's Reagent 222
Vaccination Against Typhoid Fever (A) 549	War Gases (S) 159
Vaccine, Whooping Cough (A) 543	Wars Won With Quinine Instead
Vaginal Trichomoniasis, Silver	of Artillery (A) 327
Picrate Treatment of (A) 524	Wasicky's Reagent 231
Vanilla Extract, Determination	Wassermann, Kahn and Other
of Coumarin in (A) 526	Serodiagnostic Tests for Syphi-
Vegetable Drugs, Appearance Un-	lis (A) 468
der Ultra-Violet Light 498	Water, Natural (S) 575
Venereal Disease, Control of (A) 476	Water, Public Health Aspects of
Verbena, The Holy Herb of the	the Treatment of—and Bev-
Romans (A) 322	erages With Silver (A) 141
Violamine (Ciba) 23	Water Safe for Drinking (S) 103
Virgin Forest, Six Tracts of—	Water Smelling Now a Vocation
Remain in Pennsylvania (A) 335	(A)
Virus Diseases, Chemotherapy of	Water Supplies (S) 487
(A) 521	Waxes, Chemical Microscopy of
Viruses (A) 85	Fats and 67
Viruses, The Nature of (A) 98	Weight of a Field (S) 452
	Wellcome Trust (E) 280
Vitamin A Content of Cod Liver Oil (A)	What Is Truth? (E) 536
Vitamin A Deficiency in the Eye	Wheat Germ Oil as a Valuable
(A)	Source of Vitamin E 544
Vitamin A, Isolation of Crystal-	Whooping Cough (A) 554
line (A)	Whooping Cough, The Diagnosis
Vitamin A, The Loss of—in the	of
Urine in Certain Diseases (A) 443	Whooping Cough, Pertussis En-
Vitamin A and Provitamin. Vital	dotoxin in the Treatment of (A) 554
Substances in Food and Med-	Whooping Cough, Recent Re-
icine (A) 50	ports on (A) 207
Vitamin C, Chronic Benzene Poi-	Whooping Cough Vaccine (A) 543
soning and (A) 322	Words, Words (E) 586
Vitamin C Content of Potatoes	Wound Healing and Antiseptics
(A) 317	(A) 588
Vitamin C Deficiency (A) 522	Yeasts in the Normal Mouth 167
Vitamin C, The Infra-red Ab-	Yellow Fever Threat, Health Au-
sorption Spectrum of 411	thorities Fight New (A) 332
Vitamin C, The Oxidation of (A) 188	Zinc Sulfate Pontocain Solution
Vitamin D (A) 39	Urged as a Preventive of Polio
Vitamin D, Cacao Hulls as a	(A) 518
Source of (A) 42	Zinc Sulfate and Potassium Alu-
Vitamin D, Content of Menhaden	minum Sulfate Solutions (A) 610
Fish Oil (A) 148	

PA THE

WE'RE TELLING PHYSICIANS

In Your Locality
Parke, Davis & Company of the Company

Parke, Davis & Company's systematic and extensive program of promotion with physicians is nation-wide—and so carefully planned and carried out that it reaches the doctors in the smallest towns as well as those in the largest centers.

Supplementing and supporting thousands of personal interviews every day by our large medical service staff, our constant and varied campaigns-by-mail are designed as a definite factor for bringing business to your store from those physicians who practice in your immediate vicinity.

These promotional campaigns are timed to reach the physician at regular and frequent intervals. They keep the Parke-Davis name constantly in his mind—and they all mean business, profitable business, for stores which stock Parke-Davis Products and bring them to the attention of physicians in their home neighborhoods.

For Example

Arming the Parisi-Boris Products actively armined to physicians at this time are

NATOLA. Haliver Oil fortified with Natural Vitamin D.

ABDOL with Vitamin C Five essential vitamins of standardized high potency.

GLUCO-FEDRIN . For palliative treatment of congestion of nose and throat in winter colds.

ANTUITRIN-S . . The anterior pituitary-like sex hormone.

THEELIN . . . Estrogenic hormone in chemically pure form.

PARKE, DAVIS & COMPANY . Detroit

THE WORLD'S LARGEST MAKERS OF PHARMACEUTICAL AND BIOLOGICAL PRODUCTS



Prescriptions for Profit

NLY a pharmacist can conduct a pharmacy. The prescription department is his exclusive franchise. There are other good reasons why pharmacists should exert every effort to increase prescription volume. This field of endeavor is less competitive than others. It responds to stimulation. Prescription business is profitable business.

A first essential is to have what physicians need, a

second, to stock lines and items that carry prestige, meet ready acceptance, and for which there is a created demand. Major effort can safely be expended on products bearing the Lilly Label. They qualify. They are backed by a fine reputation and they offer something definite and tangible besides low price. These are but

a few of the requisites that are met when you feature Lilly Products.

ELI LILLY AND COMPANY

INDIANAPOLIS, INDIANA, U.S.A.

Hexylresorcinol SUCRETS Promptly Soothe Irritated Throats



THE season of throat irritation is here again!

Hexylresorcinol SUCRETS displayed in this attractive new counprayed in and average ter stand are a "pick-up" item for every customer with an irritated throat condition. Dissolved slowly on the tongue, a Sucret produces a marked soothing effect on the mucous membrane of the mouth and throat . . . promptly allays that "scratchy" feeling.

Hexylresorcinol SUCRETS are convenient to carry. They are packed 24 in a neat metal box,

ped. They may be carried loose in the handbag or pocket, or kept at the bedside.

Every lot of Hexylresorcinol SUCRETS is standardized for germicidal activity by exacting bacteriological test.

Display SUCRETS in thes attractive new metal stands ... the stand comes free with an order of six dozen or more Hexylresorcinol SUCRETS. (Offer expires December

For the Conservation of Life" Pharmaceuticals - Mulford Biologicals SHARP & DOHME BALTIMORE

PHILADELPHIA

HOLADIN

An Extract of the Entire Pancreas Gland

Holadin has long proved a definite resource in pancreatic disorders, diseases associated therewith. Holadin is put up in powder in 1 oz. and ½ oz. vials, no "directions"; as commonly prescribed, the pharmacist may readily attach the usual prescription label.

FAIRCHILD BROS. & FOSTER

Holadin is also put up in capsules of 3 and 5 grains each.

New York

What are

"POPULAR SCIENCE TALKS"?

For seventeen consecutive years, the Philadelphia College of Pharmacy and Science has presented to the general public a series of talks, by members of its Faculty, on scientific subjects of popular interest.

The lectures are designed especially to combine scientific accuracy and completeness, yet with a minimum use of technical terms.

The American Journal of Pharmacy has printed many of the Popular Science Lectures of other years. Now they are assembled in book form, in 11 volumes, each volume representing an annual series.

All the books are paper-bound, and are available at the nominal cost of \$1 each, postpaid.

The books, singly or as a collection, are appropriate as holiday, birthday and graduation gifts for students, preceptors, teachers and professional men and women. Libraries of all types find them widely used as reference works, and they have met the approval of many scientific societies.

Send for a descriptive folder.

